

Rapid test for the determination of antibodies against Sars-Cov-2

July 30 2020



Stephanie Pfänder from Ruhr-Universität Bochum. Credit: RUB, Marquard

To determine immunity to SARS-Cov-2 and the effectiveness of potential vaccines, the amount of neutralizing antibodies in the blood of recovered or vaccinated individuals must be determined. A traditional



neutralization test usually takes two to three days and must be carried out with infectious coronaviruses in a laboratory complying to biosafety level 3. A Swiss-German research team from Bern and Bochum has launched a test that takes only 18 hours and doesn't have high biosafety requirements. The researchers have published their report in the journal *Vaccines* on 15 July 2020.

The test was developed at the Institute of Virology and Immunology (IVI) of the University of Bern and the Swiss Federal Office for Food Safety and Animal Health, and evaluated in cooperation with colleagues from the Ruhr-Universität Bochum (RUB) using serum samples from COVID-19 patients.

Disguising a harmless virus as SARS-Cov-2

In order to detect <u>antibodies</u> against SARS-Cov-2, the researchers used another <u>virus</u> that doesn't propagate. They exchanged the envelope protein of this virus for the spike protein of the novel coronavirus, which mediates virus entry and infection. "As a result, the viruses can be identified by antibodies against SARS-Cov-2," explains lead author Toni-Luise Meister from the Department of Molecular and Medical Virology at Ruhr-Universität Bochum. "The antibodies bind to the viruses that have been altered in this way and neutralize them so that no longer can penetrate the host cells."

Luminescence helps determine immunity

Since the virus pseudotyped in this way can't propagate in host cells, no elaborate biosafety precautions are necessary for the test. In order to determine the amount of antibodies, the researchers genetically modified the virus so that green fluorescent protein and a luciferase, an enzyme from fireflies, will be produced by infected cells. "After a single round



of infection, we can then determine how many cells show green fluorescence," says lead author Ferdinand Zettl from the Institute of Virology and Immunology in Bern. The green fluorescence is an indicator of infection with the pseudotyped virus. The less green cells the researchers are finding, the more neutralizing antibodies are present which blocked the virus. In addition, a luminometer may be used to read the luminescence signal produced by the luciferase enzyme—another way of evaluating the test.

Quick and reliable

In order to check the reliability and comparability with the conventional neutralization test, the researchers applied it to blood samples from COVID-19 patients. "The direct comparison showed a good correlation between the two test systems," explains corresponding author Professor Stephanie Pfänder from the Department of Molecular and Medical Virology at RUB. Compared to 56 hours for the conventional test, the new test is much faster, with only 18 hours to the test result. "Another great advantage is that it can be carried out in almost all medical labs, because no sophisticated safety precautions are necessary," points out Dr. Gert Zimmer from the Institute of Virology and Immunology in Bern, corresponding author of the study.

More information: Ferdinand Zettl et al. Rapid Quantification of SARS-CoV-2-Neutralizing Antibodies Using Propagation-Defective Vesicular Stomatitis Virus Pseudotypes, *Vaccines* (2020). <u>DOI:</u> 10.3390/vaccines8030386

Provided by Ruhr-Universitaet-Bochum



Citation: Rapid test for the determination of antibodies against Sars-Cov-2 (2020, July 30) retrieved 6 May 2024 from https://medicalxpress.com/news/2020-07-rapid-antibodies-sars-cov-.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.