

Scientists discover a new connection between the eyes and touch

July 6 2020



Credit: CC0 Public Domain

Tiny eye movements can be used as an index of humans' ability to anticipate relevant information in the environment independent of the information's sensory modality, a team of scientists has found. The work reveals a connection between eye movements and the sense of touch.

"The fact that tiny eye movements can hinder our ability to discriminate [tactile stimuli](#), and that the suppression of those eye movements before an anticipated tactile stimulus can enhance that same ability, may reflect that common brain areas, as well as common neural and cognitive resources, underlie both eye movements and the processing of tactile stimuli," explains Marisa Carrasco, a professor of psychology and neural science at New York University and the senior author of the paper, which appears in the latest issue of the journal *Nature Communications*.

"This connection between the eyes and touch reveals a surprising link across perception, cognition, and action," adds Stephanie Badde, an NYU post-doctoral researcher and first author of the paper.

The study asked [human participants](#) to distinguish between two kinds of vibrations ('fast'—[high frequency](#) vs. 'slow'—low frequency) that were produced by a device connected to their finger. The researchers then tracked even the tiniest of their involuntary eye movements, known as micro-saccades. These small, rapid [eye-movements](#) are known to occur even when we try to fixate our gaze on one spot. Here, participants were instructed to focus their vision on a fixation spot on a computer screen. A cue—a tap elicited by the device at their finger—would announce the next imminent vibration. What the participants did not know is that the [time interval](#) between that cue and the tactile vibration was a central part of the experimental design.

The manipulation of that interval allowed participants in some blocks to predict with more accuracy precisely when the vibration would happen. Notably, when they had that precise information, the researchers could see not only how the participants' microsaccade rates would decrease just before the vibration stimulus, but also how their ability to distinguish between fast and slow vibrations was enhanced by the suppression of micro-saccades.

More information: Stephanie Badde et al, Oculomotor freezing reflects tactile temporal expectation and aids tactile perception, *Nature Communications* (2020). [DOI: 10.1038/s41467-020-17160-1](https://doi.org/10.1038/s41467-020-17160-1)

Provided by New York University

Citation: Scientists discover a new connection between the eyes and touch (2020, July 6)
retrieved 24 April 2024 from <https://medicalxpress.com/news/2020-07-scientists-eyes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.