

SGLT-2 inhibitors may raise risk for diabetic ketoacidosis

July 29 2020



(HealthDay)—In patients with type 2 diabetes, sodium-glucose

cotransporter-2 (SGLT-2) inhibitors are associated with an increased risk for diabetic ketoacidosis (DKA), according to a study published online July 28 in the *Annals of Internal Medicine*.

Antonios Douros, M.D., Ph.D., from McGill University in Montreal, and colleagues matched 208,757 new users of SGLT-2 inhibitors with 208,757 recipients of dipeptidyl peptidase-4 (DPP-4) inhibitors to compare the risk for DKA in patients with type 2 diabetes. Data were obtained from seven Canadian provinces and the United Kingdom.

The researchers found that during 370,454 person-years of follow-up, 521 patients were diagnosed with DKA (incidence rate, 1.40 per 1,000 person-years). The risk for DKA was increased in association with SGLT-2 inhibitors versus DPP-4 inhibitors (incidence rate, 2.03 versus 0.75; hazard ratio, 2.85). Molecule-specific hazard ratios were 1.86, 2.52, and 3.58 for dapagliflozin, empagliflozin, and canagliflozin, respectively. The association was not modified by age or sex; prior insulin receipt seemed to reduce the risk.

"Because the beneficial effects of SGLT-2 inhibitors in the prevention of cardiovascular and renal disease will probably increase their uptake in the following years, physicians should be aware of DKA as a potential adverse effect," the authors write.

More information: [Abstract/Full Text \(subscription or payment may be required\)](#)

Copyright © 2020 [HealthDay](#). All rights reserved.

Citation: SGLT-2 inhibitors may raise risk for diabetic ketoacidosis (2020, July 29) retrieved 21 June 2024 from <https://medicalxpress.com/news/2020-07-sgl2-inhibitors-diabetic-ketoacidosis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.