

Significant flaw leading to underdiagnosis of gestational diabetes

July 16 2020



Credit: CC0 Public Domain

Testing procedures for gestational diabetes in pregnant women around the world need greater scrutiny to be reliable, researchers from The Australian National University (ANU) warn.

The researchers found a "significant flaw" in the approach commonly used that can lead to substantial underdiagnosis of the condition.

The problem arises if there is a delay in processing the glucose drink used to test patients' [blood samples](#). This is because the red and [white blood cells](#) continue to consume glucose. Delayed processing gives a false low glucose reading.

Lead researcher Professor Christopher Nolan, from the ANU Medical School, together with Professors Julia Potter and Peter Hickman from ACT Pathology, led a study of more than 12,000 women which found substantially higher rates of diagnosis when the [blood](#) samples were processed sooner.

Early centrifugation, within 10 minutes, compared to delayed centrifugation, increased the rate of diagnosis of gestational [diabetes](#) from 11.6 percent to 20.6 percent.

"This is a real problem leading to unreliable gestational diabetes test results in many countries around the world," Professor Nolan said.

Gestational diabetes occurs during pregnancy and can lead to increased chance of abnormally larger baby birth weights, low blood glucose levels in the newborn and [high blood pressure](#) or pre-eclampsia in the mothers.

Mothers-to-be usually undergo screening between week 24 and 28 of pregnancy and between [12-14 percent of](#) women in Australia are found to develop it.

However, the ANU researchers say that number could be significantly higher if testing protocols were improved.

Current international guidelines for testing of gestational diabetes

recommend the blood sample is placed in an ice slurry and centrifuged—where the blood sample is spun to separate the cells—within 30 minutes of collection. However, this guideline is difficult to implement, particularly if the blood collection site is distant to the laboratory, like it commonly is for rural women.

Professor Nolan said an easy fix would be to change the chemicals in the blood test collection tubes from [sodium fluoride](#) to citrate.

Sodium fluoride is used to slow down the consumption of glucose by the blood cells, but according to Professor Nolan it does a very poor job.

Citrate in the tubes, however, is much better in that it immediately stops glucose consumption, making the ice slurry and strict timing of blood processing unnecessary. The citrate stabilizes the glucose in the blood sample.

Professor Nolan says his research highlights the importance of the preanalytical processing protocol and urges that more research is needed before adopting citrate tubes widely.

In the meantime, Professor Nolan strongly encourages [pregnant women](#) to continue having [glucose](#) tolerance tests in pregnancy, as the current approach does diagnose women with [gestational diabetes](#) who are the most in need of treatment.

Provided by Australian National University

Citation: Significant flaw leading to underdiagnosis of gestational diabetes (2020, July 16) retrieved 23 April 2024 from <https://medicalxpress.com/news/2020-07-significant-flaw-underdiagnosis-gestational-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.