

# Women's egg quality dependent on metabolic factors

July 7 2020

---



Credit: CC0 Public Domain

Increasing the levels of a chemical found in all human cells could boost a woman's fertility and help select the best eggs for IVF, according to University of Queensland research.

In the world's most in-depth study of the final steps of egg maturation, the quality of a woman's eggs was found to be significantly dependent on

the important metabolic coenzyme called [nicotinamide adenine dinucleotide](#) (NAD+).

UQ Centre for Clinical Research scientist Professor Hayden Homer said NAD+ helps to ensure that eggs retain the bulk of their cellular building blocks as they mature.

"NAD+ is a critical coenzyme found in every cell in your body, and it's involved in hundreds of [metabolic processes](#), but levels decline with age," Professor Homer said.

"Egg quality declines relatively early, from the age of 30 years onwards, making it increasingly difficult to get pregnant.

"If we can maintain steady levels of NAD+ we may improve a woman's chances of getting pregnant both naturally and through IVF."

Professor Homer said an increasing number of women have to resort to IVF to have children due to delays in childbearing.

Around four per cent of all children born in Australia are the result of IVF—that's the equivalent one child in every average sized classroom.

The success rate of IVF significantly drops from 35 per cent in patients under 30 years old to just eight per cent for women over 40 years of age.

However, a quarter of Australian women undergoing IVF are over the age of 40.

Professor Homer's research team made the discovery by studying the motion of spindles, the structure that pulls chromosomes apart, in living eggs undergoing maturation.

The four-year project consisted of undertaking high-resolution time-lapse imaging of live eggs lacking the NAD<sup>+</sup> biosynthetic enzyme—Nampt.

They tracked the speed of spindles during the final stages of egg maturation and found that a "burst" of speed dependent upon NAD<sup>+</sup> is required to prevent the egg from losing too much of its building blocks.

"With [technological advances](#), this work will bring us closer to being able to select the best [eggs](#) for IVF treatment and to improving egg quality," Professor Homer said.

**More information:** Zhe Wei et al, Nampt-mediated spindle sizing secures a post-anaphase increase in spindle speed required for extreme asymmetry, *Nature Communications* (2020). [DOI: 10.1038/s41467-020-17088-6](#)

Provided by University of Queensland

Citation: Women's egg quality dependent on metabolic factors (2020, July 7) retrieved 24 April 2024 from <https://medicalxpress.com/news/2020-07-women-egg-quality-metabolic-factors.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--