

Changes in gut microbiota can greatly impact alcohol-related liver disease and cancer risk

August 27 2020



Credit: Pixabay/CC0 Public Domain

The importance of gut microbiota in alcohol-related liver disease and liver cancer has been demonstrated in two studies presented at the Digital International Liver Congress 2020. The key role of microbial biodiversity in the gut was highlighted in a study of fecal microbial transplantation. The technique shows promise as an intervention to improve some aspects of alcohol-related liver disease. A second study

used a mouse model to associate changes in gut microbiota with the action of key signaling molecules, mediating the risk of hepatocarcinogenesis.

In recent years, imbalances in gut [microbiota](#), or dysbiosis, have been implicated as contributing to alcoholic [liver](#) disease. In cases of chronic alcohol use, reactive oxygen species produced by alcohol metabolism can lead to [chronic intestinal inflammation](#), which in turn can increase gut permeability and alter microbiota composition. This includes expansion of inflammation-associated bacteria such as Proteobacteria, and reduction of protective species such as Faecalibacterium. Increased gut permeability is believed to lead to translocation of gut bacterial DNA and endotoxins to the liver. The latter, in particular, are thought to induce pro-inflammatory toll-like receptor 4 (TLR4) signaling pathways that are associated with hepatocarcinogenesis.

The importance of gut microbiota raises the possibility manipulating it to improve [patient outcomes](#). The first study tested whether fecal microbial transplant (FMT), the transfer of fecal bacteria from a healthy individual to a patient, could reduce cravings for alcohol as the first step for use in subsequent larger trials. In a double-blind, placebo-controlled, randomized pilot clinical trial, 20 patients with alcohol use disorder (AUD) and liver cirrhosis, all of whom had tried several options to quit alcohol unsuccessfully, were given FMT or placebo, with FMT shown to reduce alcohol cravings as well as total and psychosocial sickness impact profile at day 15 post-treatment. A corresponding significant increase in microbiota diversity was seen in FMT patients compared with baseline ($p=0.02$), including a higher relative abundance of *Odoribacter*. *Alistipes* and *Roseburia* were also more abundant in patients given FMT compared with placebo at day 15.

"FMT was safe and is shown to have an impact on reducing short-term alcohol cravings and improving psychosocial quality of life in patients

with cirrhosis and AUD," said study presenter Dr. Jasmohan S Bajaj of McGuire VA Medical Center, U.S. "The relative abundance of short-chain-fatty-acid-producing bacteria identified in patients with higher diversity after FMT demonstrates that altering the gut-brain axis is a potential avenue to alleviating AUD in those with cirrhosis."

A second study explored how [gut microbiota](#) may affect the process of developing hepatocellular carcinoma. The study used mice genetically engineered to develop steatohepatitis (NEMO-hepa mice). By crossing these mice with others with inactivated genes involved in the inflammatory response to bacteria, and then altering the gut microbial balance with broad-spectrum antibiotics, the research team showed that knocking out the NLRP6 receptor (a key mediator of colonic homeostasis that can cause intestinal dysbiosis if deficient) leads to more severe steatohepatitis and a higher tumor burden. The degree of intestinal barrier permeability was highly correlated with tumor burden as well as several indicators of inflammation in the liver. Crucially, this immune phenotype could be transferred to other mice by FMT, provided they had functional TLR4 signaling, and could be reversed if the transplanted microbiota were depleted with [broad-spectrum antibiotics](#).

"Strikingly, we also found that replacing depleted Akkermansia muciniphila bacteria in the guts of these mice helped ameliorate their inflammation and steatohepatitis," said Dr. Kai Markus Schneider of University Hospital RWTH Aachen, Germany. "This knowledge of how short-term changes to microbiota reshape the hepatic tumor microenvironment has the potential to reveal new therapeutic options for cancer prevention and therapy."

"The understanding of interactions between the human and microbiome genome (metagenome) in health and disease has represented one of the major areas of progress in the last few years," said Professor Luca Valenti, an EASL Scientific Committee member from the University of

Milan, Italy. "These studies lay the groundwork for exploiting this new knowledge for the treatment of liver disease."

More information: Digital International Liver Congress 2020: ilc-congress.eu/

Provided by Spink Health

Citation: Changes in gut microbiota can greatly impact alcohol-related liver disease and cancer risk (2020, August 27) retrieved 27 April 2024 from <https://medicalxpress.com/news/2020-08-gut-microbiota-greatly-impact-alcohol-related.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.