

High fructose diet in pregnancy impacts metabolism of offspring, study finds

August 18 2020, by Cheryl Norrie



Credit: CC0 Public Domain

An increased level of fructose intake during pregnancy can cause significant changes in maternal metabolic function and milk composition and alter the metabolism of their offspring, researchers from the

University of Otago, Wellington, have found.

The research, which was led by Dr. Clint Gray, a Research Fellow in the Department of Pediatrics and Child Health, found increasing the fructose in the diets of female guinea pigs led to highly significant and consistent changes in the [free fatty acids](#) circulating in the blood of their offspring. This was despite the offspring consuming no fructose themselves.

The research is published in the international journal *Frontiers in Endocrinology*.

First author, Ph.D. student Erin Smith, says "previous research has shown poor quality nutrition during pregnancy can predispose offspring to long-term consequences, including the development of obesity, diabetes and cardiovascular disease later in life. However, there has been a lack of data examining the impact of increased fructose intake before and during pregnancy and subsequent adverse effects on lactation, fetal development and offspring metabolic function."

The two [experimental groups](#) were fed either a control [diet](#) or a fructose diet prior to and during pregnancy. The fructose group was given supplementary fructose water to replicate increased sugar-sweetened beverage intake 60 days prior to mating and until the delivery of their offspring. Fructose made up 16.5 percent of their diets, closely resembling the average human consumption of fructose/sugar in Western countries, which is estimated at about 14 percent of average daily caloric intake.

"We found fructose had a significant impact on a pregnant females' metabolic status and the free fatty acid content of their milk. We also provide the first evidence that [offspring](#) born from fructose-fed mothers display a very specific pattern of increased free fatty acids and altered

[lipid metabolism](#) that persists throughout early life."

Ms Smith says it is well known that increased levels of circulating free fatty acids increases the risk of obesity, insulin resistance, type 2 diabetes and [cardiovascular disease](#)—with increased fatty acid synthesis shown to occur following fructose consumption.

She says the evidence suggests suboptimal maternal diets, such as diets high in fructose and refined sugars, may be contributing to the rise in metabolic diseases in humans observed during the past 40 to 50 years.

"Our study emphasizes the importance of limiting added refined [fructose](#), such as sugar-sweetened beverages, and striving for a more nutritionally balanced diet in women prior to and during pregnancy and lactation."

The article, "Fructose Consumption During Pregnancy Influences Milk Lipid Composition and Offspring Lipid Profiles in Guinea Pigs" is published in the international journal *Frontiers in Endocrinology*.

More information: Erin Vanessa LaRae Smith et al. Fructose Consumption During Pregnancy Influences Milk Lipid Composition and Offspring Lipid Profiles in Guinea Pigs, *Frontiers in Endocrinology* (2020). [DOI: 10.3389/fendo.2020.00550](https://doi.org/10.3389/fendo.2020.00550)

Provided by University of Otago

Citation: High fructose diet in pregnancy impacts metabolism of offspring, study finds (2020, August 18) retrieved 7 May 2024 from <https://medicalxpress.com/news/2020-08-high-fructose-diet-pregnancy-impacts.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.