

Nicotine withdrawal in zebrafish offers clues to human addiction

August 21 2020, by Adrianna MacPherson



Credit: Pixabay/CC0 Public Domain

Researchers have identified nicotine-induced withdrawal for the first time in the zebrafish model—a discovery that could be used to test ways of treating withdrawal from addiction in humans.



Trevor Hamilton, an adjunct professor in the Faculty of Science and member of the Neuroscience and Mental Health Institute at the University of Alberta, examined the role of nicotine exposure on zebrafish. He looked at two main factors: their <u>boldness</u> and their locomotion.

With acute nicotine exposure, the zebrafish demonstrated increased boldness, as measured by the fish's inclination to approach a small figurine placed in the testing arena. Acute exposure also increased the fish's locomotion, or willingness to move around the tank.

Repeated nicotine exposure followed by a period of <u>withdrawal</u>, however, caused a decrease in boldness and movement, which from previous research is thought to be associated with increased anxiety-like behavior in the zebrafish.

Hamilton linked the traits to emotionality during addiction in humans: "We know that people who quit smoking have an increase in anxiety, and that increased feeling of stress drives them to go have another cigarette. That's part of the addictive mechanism. So at a very basic level we're looking at the same thing in fish."

Hamilton explained that zebrafish make an ideal model for pharmacological research because their genome has been sequenced and they have a rapid reproduction rate, which allows researchers to easily examine patterns of addiction to substances like <u>nicotine</u> across generations and with powerful genetic tools.

The next steps in the project involve narrowing the window of time to determine when the shift from an increase to decrease in boldness occurs, and pinpointing the start of the withdrawal behavior.

"If we can find that point and then also at the same time look before and



after at changes in <u>brain chemistry</u>, it'll give us an idea of which neurotransmitters are involved, and where neuroadaptation is taking place," said Hamilton.

"Then, once we learn a little bit more about the brain chemistry, we can start looking at which drugs could potentially alleviate and somewhat reverse these adaptations."

The study, "Opposing Effects of Acute and Repeated Nicotine Exposure on Boldness in Zebrafish," was published in *Scientific Reports*.

More information: Rachel Dean et al. Opposing effects of acute and repeated nicotine exposure on boldness in zebrafish, *Scientific Reports* (2020). DOI: 10.1038/s41598-020-65382-6

Provided by University of Alberta

Citation: Nicotine withdrawal in zebrafish offers clues to human addiction (2020, August 21) retrieved 5 May 2024 from https://medicalxpress.com/news/2020-08-nicotine-zebrafish-clues-human-addiction.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.