

No safe level of caffeine consumption for pregnant women and would-be mothers

August 24 2020



Credit: CC0 Public Domain

Women who are pregnant or trying to conceive should be advised to avoid caffeine because the evidence suggests that maternal caffeine consumption is associated with negative pregnancy outcomes and that

there is no safe level of consumption, finds an analysis of observational studies published in *BMJ Evidence Based Medicine*.

Caffeine is probably the most widely consumed psychoactive substance in history, and many people, including pregnant women consume it on a daily basis.

Pregnant women have been advised that consuming a small amount of caffeine daily will not harm their baby. The UK NHS, the American College of Obstetricians and Gynecologists, the Dietary Guidelines for Americans and the European Food Safety Authority (EFSA) set this level at 200 mg caffeine, which approximates to roughly two cups of moderate-strength coffee per day.

This study undertook a review of current evidence on caffeine-related [pregnancy outcomes](#), to determine whether the recommended safe level of consumption for pregnant women is soundly based.

Through database searches, Professor Jack James, of Reykjavik University, Iceland, identified 1,261 English language peer-reviewed articles linking caffeine and caffeinated beverages to [pregnancy outcomes](#).

These were whittled down to 48 original observational studies and meta-analyses published in the past two decades reporting results for one or more of six major negative pregnancy outcomes: miscarriage, stillbirth, low birth weight and/or small for gestational age, [preterm birth](#), childhood acute leukaemia, and childhood overweight and obesity.

A total of 42 separate findings were reported in 37 observational studies; of these 32 found that caffeine significantly increased risk of adverse pregnancy outcomes and 10 found no or inconclusive associations. Caffeine-related risk was reported with moderate to high levels of

consistency for all pregnancy outcomes except preterm birth.

Eleven studies reported on the findings of 17 meta-analyses, and in 14 of these maternal caffeine consumption was associated with increased risk for four adverse outcomes: miscarriage, stillbirth, low birth weight and/or small for gestational age, and childhood acute leukaemia. The three remaining meta-analyses did not find an association between maternal caffeine consumption and preterm birth.

No meta-analyses looked at the association between maternal caffeine consumption and childhood overweight and obesity, but four of five [observational studies](#) reported significant associations.

This is an observational study, so can't establish causation, and the author points out that the results could be impacted by other confounding factors, such as recall of caffeine consumption, maternal cigarette smoking and most importantly pregnancy symptoms. Pregnancy symptoms such as nausea and vomiting in early pregnancy are predictive of a healthy pregnancy and women who experience them are likely to reduce their caffeine intake.

But he adds that the dose-responsive nature of the associations between caffeine and adverse pregnancy outcomes, and the fact some studies found no threshold below which negative outcomes were absent, supports likely causation rather than mere association.

Professor James concludes that there is "substantial cumulative evidence" of an association between maternal caffeine consumption and diverse negative pregnancy outcomes, specifically miscarriage, stillbirth, [low birth weight](#) and/or small for gestational age, childhood acute leukaemia and childhood overweight and obesity, but not preterm birth.

As a result, he adds, current health recommendations concerning

caffeine consumption during pregnancy are in need of "radical revision."

"Specifically, the cumulative scientific evidence supports [pregnant women](#) and women contemplating pregnancy being advised to avoid [caffeine](#)," he says.

More information: Maternal caffeine consumption and pregnancy outcomes: a narrative review with implications for advice to mothers and mothers-to-be, *BMJ Evidence Based Medicine*, [DOI: 10.1136/bmjebm-2020-111432](#)

Provided by British Medical Journal

Citation: No safe level of caffeine consumption for pregnant women and would-be mothers (2020, August 24) retrieved 24 May 2024 from <https://medicalxpress.com/news/2020-08-safe-caffeine-consumption-pregnant-women.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--