

In study, acetaminophen makes risky moves seem less dangerous

September 8 2020, by Jeff Grabmeier



Credit: Pixabay/CC0 Public Domain

While acetaminophen is helping you deal with your headache, it may also be making you more willing to take risks, a new study suggests.

People who took [acetaminophen](#) rated activities like "bungee jumping off a tall bridge" and "speaking your mind about an unpopular issue in a meeting at work" as less risky than people who took a placebo, researchers found.

Use of the drug also led people to take more risks in an experiment where they could earn rewards by inflating a virtual balloon on a computer: Sometimes they went too far and the balloon popped.

"Acetaminophen seems to make people feel less negative emotion when they consider risky activities—they just don't feel as scared," said Baldwin Way, co-author of the study and associate professor of psychology at The Ohio State University.

"With nearly 25 percent of the population in the U.S. taking acetaminophen each week, reduced risk perceptions and increased risk-taking could have important effects on society."

The study extends a series of studies led by Way that have shown acetaminophen—the main ingredient in the pain-reliever Tylenol and nearly 600 other medicines—has [psychological effects](#) that most people don't consider when they take it.

Previous research by Way and his colleagues has shown that acetaminophen reduces positive and [negative emotions](#), including hurt feelings, distress over another's suffering and even your own joy.

Way conducted the current study with Alexis Keaveney, a former doctoral student at Ohio State, and Ellen Peters, a former professor at Ohio State who is now at the University of Oregon. The study was published online in the journal *Social Cognitive and Affective Neuroscience*.

In one study, 189 [college students](#) came to a lab and took either 1,000 mg of acetaminophen (the recommended dosage for a headache) or a placebo that looked the same. After waiting for the drug to take effect, the participants rated on a scale of 1 to 7 how risky they thought various activities would be.

Results showed that those under the influence of acetaminophen rated activities like bungee jumping, walking home alone at night in an unsafe area of town, starting a new career in your mid-30s, and taking a skydiving class as less risky than those who took the placebo.

The effects of acetaminophen on risk-taking were also tested in three separate experimental studies.

Across these studies, 545 undergraduate students took part in a task developed in 2002 that is often used by researchers to measure risk-taking behavior. Other researchers have shown that taking more risk on this task predicted risky behaviors outside the laboratory, including alcohol and drug use, driving without a seatbelt and stealing.

In the task, participants click a button on the computer to inflate a balloon on their computer screen. Each time they inflate it they receive virtual money. They can stop at any time and add the money to their "bank," and move on to the next balloon. But there is risk involved.

"As you're pumping the balloon, it is getting bigger and bigger on your computer screen, and you're earning more money with each pump," Way said.

"But as it gets bigger you have this decision to make: Should I keep pumping and see if I can make more money, knowing that if it bursts I lose the money I had made with that balloon?"

For those who took the acetaminophen, the answer was: Keep on pumping. Results showed that those on the drug pumped more times than those on the placebo and had more burst balloons.

"If you're risk-averse, you may pump a few times and then decide to cash out because you don't want the balloon to burst and lose your money," he said.

"But for those who are on acetaminophen, as the balloon gets bigger, we believe they have less anxiety and less negative emotion about how big the [balloon](#) is getting and the possibility of it bursting."

The results have a variety of real-life implications, Way said.

For example, acetaminophen is the recommended treatment by the CDC for initial COVID-19 symptoms.

"Perhaps someone with mild COVID-19 symptoms may not think it is as risky to leave their house and meet with people if they're taking acetaminophen," Way said.

Even everyday activities like driving presents people with constant decisions involving risk perception and assessment that could be altered by use of the painkiller.

"We really need more research on the effects of acetaminophen and other over-the-counter drugs on the choices and risks we take," he said.

More information: Alexis Keaveney et al. Effects of acetaminophen on risk taking, *Social Cognitive and Affective Neuroscience* (2020). [DOI: 10.1093/scan/nsaa108](https://doi.org/10.1093/scan/nsaa108)

Provided by The Ohio State University

Citation: In study, acetaminophen makes risky moves seem less dangerous (2020, September 8) retrieved 20 June 2024 from <https://medicalxpress.com/news/2020-09-acetaminophen-risky-dangerous.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.