

Antacid monotherapy more effective in relieving epigastric pain than in combination with lidocain

September 29 2020

Antacid monotherapy is more effective in relieving epigastric pain than in combination with lidocaine. That is the conclusion of a study to be published in the September 2020 issue of *Academic Emergency Medicine* (AEM), a journal of the Society for Academic Emergency Medicine (SAEM).

The lead author of the single-center study is Dr. Jaimee Warren, a first-year physician at the Royal Melbourne Hospital, Parkville, Victoria, Australia. The findings of the study are discussed with the author in a recent AEM podcast, [We Didn't Start the Fire But Can Antacid Monotherapy Stop the Fire?](#)

The double-blind, randomized clinical trial compared three different solutions for the treatment of adults with epigastric pain or dyspepsia presenting to the [emergency department](#) (ED): antacid [monotherapy](#), antacid/lidocaine 2% solution, and antacid/lidocaine 2% viscous gel.

Warren, et al. concluded that all three treatments worked and there was no statistical difference in [pain relief](#) among the groups at 30 and 60 minutes; however, antacid monotherapy was found to be the most palatable solution, with statistically significant differences in taste, bitterness, and overall acceptability, and there were fewer side effects. In conclusion, the study authors recommend using antacid monotherapy in place of lidocaine/antacid combination therapy for management of

dyspepsia and epigastric pain for patients in the ED.

Commenting on the study is Robert Ehrman, MD, MS, research faculty in the department of emergency medicine at Wayne State School of Medicine in Detroit, MI:

"This study nicely demonstrates that 'more' is not always synonymous with 'better' when it comes to treating pain. Antacid alone appears to be equally efficacious to [antacid](#)-lidocaine combinations, without any of the unpleasant side effects that may discourage its use. This is important information to provide to patients as antacids are available over-the-counter, thereby allowing patients to manage their symptoms without a visit to the emergency department."

More information: Jaimee Warren et al, Antacid Monotherapy Is More Effective in Relieving Epigastric Pain Than in Combination With Lidocaine: A Randomized Double-blind Clinical Trial, *Academic Emergency Medicine* (2020). [DOI: 10.1111/acem.14069](https://doi.org/10.1111/acem.14069)

Provided by Society for Academic Emergency Medicine

Citation: Antacid monotherapy more effective in relieving epigastric pain than in combination with lidocain (2020, September 29) retrieved 26 April 2024 from <https://medicalxpress.com/news/2020-09-antacid-monotherapy-effective-relieving-epigastric.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.