


Bacteria virus combo may be cause of neonatal brain infections in Uganda

September 30 2020



An international team led by Penn State investigated the origin of the severe brain infections afflicting thousands of Ugandan infants in the neonatal period, and often resulting in hydrocephalus in the survivors. Fluid from the brain taken at surgery led to the recovery of a novel strain of a highly virulent bacteria, shown on the tree of life of related organisms. These infection cases, shown as red dots on the map, are concentrated within a region in Eastern Uganda characterized by wetlands and swamps on the north and south banks of Lake Kyoga that the Nile river flows into and out of. This map is adapted and based on OCHA/ReliefWeb (UN Office for the Coordination of Humanitarian Affairs) and is used as permitted by the agency. Credit: Schiff Lab, Penn State

A newly identified bacteria and a common virus may be the underlying cause of infection-induced hydrocephalus in Uganda, according to an international team of researchers.

"Thirteen years ago, while visiting Uganda and seeing a stream of kids with hydrocephalus after [infection](#) I asked the doctors, 'What is the biggest problem you have that you can't solve?'" said Steven J. Schiff, Brush Chair Professor of Engineering and professor of engineering science and mechanics, neurosurgery and physics, Penn State. "'Why don't you figure out what makes these kids sick?' was the reply."

By that time, the doctors at CURE Children's Hospital of Uganda, had seen more than a thousand infants with infection-caused hydrocephalus and were unable to culture a single thing in the laboratory. They have now seen over 8,000 similar children in this one, small Ugandan hospital.


"Hydrocephalus is the most common childhood neurosurgical condition that we see in the population that we serve," said Edith Mbabazi-Kabachelor, director of research, CURE Children's Hospital of Uganda. "If hydrocephalus is left untreated in children less than two years old, the progressive increase in head size will lead to further brain damage, resulting in the majority of these children dying, and those that survive being left with severe cognitive and physical disability."

Severe systemic bacterial infection during the first four weeks of life accounts for an estimated 680,000 to 750,000 yearly neonatal deaths worldwide. Hydrocephalus is the most common brain disorder in childhood and the largest single cause of childhood hydrocephalus is neonatal infection, accounting for an estimated 160,000 yearly cases, said Schiff.

Over a 5-year study in Uganda, supported by the U.S. National Institutes of Health, using advanced genomic techniques the team uncovered the

major bacterial and viral underpinnings of these infections, the researchers report today (Sept. 30) in *Science Translational Medicine*.

Schiff and his team have studied this problem for more than 10 years, but in the last five years, they took a different approach, using DNA and RNA sequencing techniques to identify the causative agents. The researchers looked at blood and [cerebrospinal fluid](#) drawn from 100 cases of post-infectious hydrocephalus and control patients without infection in Uganda. There were 64 infants with post-infectious hydrocephalus and 36 with non-post-infectious hydrocephalus. All infants were under three months old. The researchers prepared the samples in two ways—fresh-frozen and preserved—and they sent samples to two different laboratories in the U.S., where samples were analyzed with different techniques. This was to ensure valid and reproducible results.



CT brain images from patients with post-infectious hydrocephalus (PIH), non-postinfectious hydrocephalus (NPIH), *Paenibacillus* infection (Paeni), or cytomegalovirus infection (CMV). Credit: J.N. Paulson et al., *Science Translational Medicine* (2020)

"We found this weird bacteria dominating," said Schiff.

The bacteria was a previously unidentified strain of *Paenibacillus thiaminolyticus*, now named *Mbale* after the city where the CURE

Children's Hospital is located.

"The initial link between hydrocephalus and *Paenibacillus* was made through high-throughput sequencing and PCR analyses at the Center for Infection and Immunity in the Mailman School of Public Health at Columbia University, a renowned center led by W. Ian Lipkin," said Schiff.

High-throughput sequencing allows sequencing of more than one DNA molecule at the same time, and PCR analysis multiplies existing DNA samples so that they are easier to analyze and identify.

"You build a field of dreams—in this case a platform for pathogen discovery—and wait for the right partner and the right project," said W. Ian Lipkin, John Snow Professor of Epidemiology and director, Center for Infection and Immunity, Mailman School of Public Health, and professor of pathology and neurology, College of Physicians & Surgeons, Columbia University. "Steven Schiff is a remarkable investigator and this is such a project. It stands out for impact amongst hundreds we've done over a period of more than 30 years. Our team is delighted to have had an opportunity to help implicate an agent and contribute to control of this devastating disorder."

The researchers managed to grow the difficult-to-culture new bacterial strain at Penn State, and tested it on mice. While the common variants of *Paenibacillus* are harmless, the Mbale strain was lethal to the mice.

The researchers found the new bacterial strain in the cerebrospinal fluid of the infection-induced hydrocephalic children and then only in the youngest patients.

"While we tested infants up to three months old, we mostly identified the cause of infections in those less than six weeks of age," said Schiff. "If

we didn't study them really early in life, then the infection had already burned out. Between 6 and 12 weeks there were very few positive results."

Schiff was not satisfied with finding the proposed bacterial cause of the problem, he said. He reasoned that other diseases had both a bacterial and viral component and so the team looked for viral, fungal and parasitic genetic material. They found cytomegalovirus (CMV) in the cerebrospinal fluid of the infection-caused hydrocephalic infants, but not in that of the other hydrocephalus patients.

CMV is a common virus found around the world. The virus causes minor symptoms, if any, in most adults, although babies may be born with congenital CMV or acquire it early in life and be significantly harmed by neurological damage. The researchers only found CMV in the cerebrospinal fluid of babies with post-infection hydrocephalus.

While the researchers believe they have found the source of the infections that cause the high prevalence of hydrocephalus, they do not know where the babies encounter the new bacteria. According to Schiff, the bacteria may be soil- or water-born and more work is necessary to find the bacterial source.

The researchers are creating predictive models that, coupled with data they are now analyzing from thousands of infants and satellite-acquired rainfall to predict optimal treatment for individual locations. The researchers said they do not know if this particular bacterial virus combination exists outside this area of Uganda. However, the same strategy of using DNA and RNA to diagnose previously unknown causes of similar infections can be used in many other regions in the developing world where similar cases are seen.

"Both care and prevention fit within an economic framework that will

help plan the optimal apportionment of health care funds between treatment and prevention, so that the health of these infants is best addressed," said Mbabazi-Kabachelor. "These strategies will help guide optimal use of resources for better long-term patient outcomes."

More information: J.N. Paulson at Genentech Inc. in South San Francisco, CA et al., "Paenibacillus infection with frequent viral coinfection contributes to postinfectious hydrocephalus in Ugandan infants," *Science Translational Medicine* (2020).

stm.sciencemag.org/lookup/doi/10.1126/scitranslmed.aba0565

Provided by Pennsylvania State University

Citation: Bacteria virus combo may be cause of neonatal brain infections in Uganda (2020, September 30) retrieved 2 May 2024 from <https://medicalxpress.com/news/2020-09-bacteria-virus-combo-neonatal-brain.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--