

Children with social anxiety, maternal history of depression more likely to develop depression

September 3 2020



Credit: CC0 Public Domain

Although researchers have known for decades that depression runs in families, new research from Binghamton University, State University of



New York, suggests that children suffering from social anxiety may be at particular risk for depression in the future.

"We already know from previous research that <u>children</u> with social anxiety symptoms are at high risk of developing depression, as are offspring of depressed mothers," said Holly Kobezak, co-author of the new paper and lab manager at the Mood Disorders Institute at Binghamton University. "Our findings take what is already known one step further by suggesting that the combination of these risk factors may be even more insidious than the presence of either risk factor alone."

The researchers invited approximately 250 eight- to 14-year-old children whose mothers either did or did not have a history of major depressive disorder (MDD) into their lab to complete questionnaires measuring social anxiety and depression symptoms. Symptoms were reassessed at six-month intervals over a period of two years in order to capture changes in <u>symptom</u> levels over time as children progressed further into adolescence, which is a critical time period for the development of depression.

Their results showed that high levels of social anxiety predicted increases in <u>depression symptoms</u> over time, but only among children of mothers with a history of MDD.

"This provides preliminary evidence that risk for the development of depression among children with social anxiety may be particularly high among children who are already at risk for depression based on a maternal history of the disorder," Kobezak said.

According to Kobezak, these findings allow the researchers to pinpoint a subgroup of children who may be at particularly high risk for developing depression on the basis of two established risk factors that have rarely, if ever, been evaluated in combination.



"This information is useful because it can help us more precisely identify children at need of early intervention and may lay the groundwork for research that works to identify mechanisms of risk that can be targeted in clinical interventions for this group of children," she said.

The researchers hope that their findings will encourage others to explore the specific ways in which social <u>anxiety</u> symptoms and exposure to maternal depression may work together to increase risk of depression in children over time.

"An important point is that our findings provide insight into the circumstances that may put children at heightened risk of depression, but equally important is research that will help us understand why this may be true," said Kobezak. "With that said, we hope future research will investigate additional variables that can explain the impact of the transactional relationship between social anxiety and maternal MDD on depression. For example, future research could focus on disruptions in social functioning and interpersonal relations resulting from these experiences and whether this might be why these children are at such elevated risk. If so, this could be specifically targeted by interventions."

More information: Holly M. Kobezak et al, Prospective associations between social anxiety and depression in youth: The moderating role of maternal major depressive disorder, *Journal of Adolescence* (2020). DOI: 10.1016/j.adolescence.2020.05.003

Provided by Binghamton University

Citation: Children with social anxiety, maternal history of depression more likely to develop depression (2020, September 3) retrieved 1 May 2024 from <u>https://medicalxpress.com/news/2020-09-children-social-anxiety-maternal-history.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.