


Cigarette smoking associated with worse outcomes for bladder cancer patients after surgery

September 15 2020



Credit: CC0 Public Domain

Patients treated for bladder cancer with a surgery known as radical cystectomy have worse outcomes if they are smokers, according to a systematic review and meta-analysis by Keck Medicine of USC. The

study appeared in *The Journal of Urology*.

"This study is important because while it is known that tobacco smoking is the leading cause of [bladder cancer](#), this is the first study to suggest that smoking puts [bladder](#) cancer patients at risk after diagnosis," says Giovanni Cacciamani, MD, lead author of the study and assistant professor of research urology at the Keck School of Medicine of USC.

More than 500,000 cases of bladder cancer are diagnosed each year worldwide. When the cancer is large or has spread beyond the bladder, patients are typically treated with chemotherapy followed by a radical cystectomy, which is when the bladder, nearby lymph nodes and surrounding organs are surgically removed.

Cacciamani and fellow Keck Medicine researchers searched databases to select 17 studies that reported on the impact of [tobacco smoking](#) on chemotherapy response and survival outcomes of 13,777 patients following radical cystectomy. Of these patients, 40.8% were active [smokers](#) at the time of the surgery, 14.1% former smokers and 45.1% had never smoked or were not smoking at the time of the surgery.

The study showed that active smokers responded worse to chemotherapy and had higher mortality rates, both in general and specifically from bladder cancer, and a higher rate of bladder cancer recurrence than patients who never smoked or were not smoking at the time of surgery.

Former smokers also fared worse in these categories than those who had never smoked, even though the differences were less significant.

One reason smoking affects outcomes is that nicotine has shown to suppress the body's immune system, leading to more complications, according to the study authors. "In addition, patients with a history of smoking tend to have more aggressive forms of cancer and, if they

survive bladder cancer, are more at risk for other potentially fatal cancers, such as lung cancer," says Cacciamani.

Looking forward, the researchers recommend that [health care professionals](#) advise bladder cancer patients to stop smoking after their diagnoses. "The research suggests that as long as a person is not smoking at the time of chemotherapy and surgery, they might do better," says Cacciamani.

He also recommends that physicians monitor smokers more carefully post-surgery than other patients because they are more at risk for complications or death.

In addition, the study authors recommend that future studies or [clinical trials](#) involving bladder cancer chart patients' smoking status to create a more accurate picture of what factors affect [cancer](#) survival and recurrence.

"While this study does not answer all the questions, it is an excellent starting point for investigating the association between [smoking](#) and long-term oncological outcomes, and will hopefully lead to new protocols that will benefit patients," says Cacciamani.

More information: G. E. Cacciamani et al, Association between Smoking Exposure, Neoadjuvant Chemotherapy Response and Survival Outcomes following Radical Cystectomy: Systematic Review and Meta-Analysis, *Journal of Urology* (2020). [DOI: 10.1097/JU.0000000000000813](#)

Provided by University of Southern California

Citation: Cigarette smoking associated with worse outcomes for bladder cancer patients after surgery (2020, September 15) retrieved 10 April 2024 from <https://medicalxpress.com/news/2020-09-cigarette-worse-outcomes-bladder-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.