

Combining tests at point of care dramatically increases COVID-19 detection in hospitalised patients

September 2 2020, by Craig Brierley



Man taking COVID-19 test. Credit: [U.S. Pacific Fleet](#)

A Cambridge hospital has piloted the use of combined rapid point-of-care nucleic acid and antibody testing for SARS-CoV-2 infection after researchers at the University of Cambridge showed that this approach was superior to virus detection alone for diagnosing COVID-19 disease.

Point-of-care testing—in other words, testing patients as soon as they

arrive at the hospital—is essential for enabling healthcare workers to rapidly diagnose patients and direct those who test positive for infection to dedicated wards. A recent study showed that SAMBA II, a new point-of-care PCR test for SARS-CoV-2 developed by Cambridge researchers, was able to dramatically reduce time spent on COVID-19 "holding" wards—allowing patients to be treated or discharged far quicker than with current lab testing set-ups.

PCR tests involve extracting a miniscule amount of RNA from the virus and copying it millions of times, creating an amount large enough to confirm presence of the virus. The virus is captured through a swab inside the nostrils and at the back of the throat. However, it can take as long as 14 days for an individual to show symptoms of COVID-19, by which time the virus may have moved away from the nose and throat and into the lungs and other tissues and organs, making it harder to detect via a swab test. As a result, studies have shown that PCR tests can miss as many as a half of infected patients five days after infection.

Antibody tests provide an alternative way of identifying infected individuals, but [antibodies](#)—molecules produced by our immune system in response to infection—generally do not appear until at least six days after infection.

Professor Ravi Gupta from the Cambridge Institute of Therapeutic Immunology and Infectious Disease at the University of Cambridge said: "We still do not have a gold standard test for diagnosing COVID-19. This poses a challenge to healthcare workers who need to make quick and safe decisions about how and where to treat patients. The two main types of test—PCR and antibody tests—both have limitations because of the nature of [coronavirus](#) infection and how our body responds. But we've shown that if you combine them and carry out both at point of care, their reliability can be hugely increased."

Professor Gupta led a team that used the approach of combining rapid point-of-care PCR and antibody tests to diagnose 45 patients at Addenbrooke's Hospital, Cambridge University Hospitals NHS Foundation Trust. The results of this peer-reviewed study are published in *Cell Reports Medicine*.

The patients, each of whom had suspected moderate to severe COVID-19 disease, provided nose/throat swabs for the tests detecting nucleic acid (virus genetic material) and blood serum for antibody testing an average (median) of seven days after the onset of illness.

The authors designed a gold standard reference test made of two parts, either of which could be positive to confirm COVID-19. The first part was an in vitro test where artificial SARS-CoV-2 viruses were made and mixed with serum from patients to see whether the serum contained neutralizing antibodies. The second part of the gold standard was the standard Public Health England laboratory test looking for genetic viral material in nose/throat swabs. Using this gold standard, 24 of the patients had COVID-19.

Professor Gupta's team used SAMBA II machines, developed by Cambridge spinout company Diagnostics for the Real World, for the nucleic acid tests, and a combination of two finger prick antibody tests, both of which test for antibodies against the spike protein on the surface of the SARS-CoV-2 virus.

Overall, the nucleic acid tests could identify eight out of ten patients with COVID-19, but when combined with the rapid antibody tests, 100% of the COVID-19 patients were correctly identified. Among the 21 patients who did not have COVID-19, there were four false positive results with one antibody test and only one false positive with the second antibody test, demonstrating that one performed better than the other.

"Combining point-of-care PCR and antibody testing could be a game-changer for rapidly identifying those patients with moderate to severe COVID-19 infection," said Professor Gupta. "This could prove extremely useful, particularly in the event of a second wave arising during flu season, when it will not be immediately clear whether the patients had COVID-19 or seasonal flu."

Professor Gupta envisages that hospitals deploying this approach would carry out a finger prick blood test and nose/throat swab at the same time on admission to hospital. The antibody test result is available within 15 minutes, but might benefit from confirmation with a second point-of-care antibody test. Importantly the study showed that the antibody tests can detect antibodies against a mutated form of SARS-CoV-2, D614G in spike protein, that has now become the dominant strain worldwide.

This approach could be particularly beneficial in low resource settings where centralized virology laboratories are scarce and the pandemic is expanding, said Professor Gupta. In addition, it removes the need for repeated nose/throat swabbing when the first [test](#) is negative and suspicion of COVID-19 is high, which may generate aerosols and lead to transmission.

More information: Petra Mlcochova et al. Combined point of care nucleic acid and antibody testing for SARS-CoV-2 following emergence of D614G Spike Variant, *Cell Reports Medicine* (2020). [DOI: 10.1016/j.xcrm.2020.100099](https://doi.org/10.1016/j.xcrm.2020.100099)

Provided by University of Cambridge

Citation: Combining tests at point of care dramatically increases COVID-19 detection in hospitalised patients (2020, September 2) retrieved 3 May 2024 from

<https://medicalxpress.com/news/2020-09-combining-covid-hospitalised-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.