

## Hypercoagulability in patients with COVID-19

September 24 2020



A, 76-year-old woman who presented to emergency department with upper back pain. Axial abdominopelvic CT angiogram shows nonocclusive acute aortic thrombus (arrow). Bilateral common iliac artery and left internal iliac artery thrombi (not shown) were also seen, as well as signs of chronic atherosclerotic disease. After findings of concurrently performed chest CT raised concerns, patient was determined to have COVID-19. Credit: American Roentgen Ray Society (ARRS), American Journal of Roentgenology (AJR)



According to an open-access article in ARRS' *American Journal of Roentgenology* (AJR), because coronavirus disease (COVID-19) may present with medium to large arterial and venous abdominal and extremity macrothrombi, radiologists should raise concern for COVID-19 when identifying thromboembolic abdominopelvic findings.

Bari Dane of NYU Langone Medical Center in New York City and colleagues at NYU Winthrop in Mineola, New York retrospectively compared 82 <u>patients</u> (58 men, 24 women; mean age 58.8) with COVID-19 who underwent abdominal ultrasound or CT with 82 patients (43 men, 39 women; mean age 53.5) without COVID-19 for thromboembolism and solid-organ infarction.

Nine (11%) patients with COVID-19 had thromboembolic findings (seven men, two women; mean age 67.0). Thromboembolism was described in four of the 38 CT reports (10.5%). Two others described splenic infarctions with patent vasculature (overall positivity rate, 15.8% [6/38]). Three patients had arterial thromboembolism in the aorta or major branches.

Noting one patient without COVID-19 had known portal vein thrombus on CT, "thromboembolic findings occurred more frequently in patients with than without COVID-19 (p = 0.02)," Dane et al. concluded.

Acknowledging this study's small sample size and single-institution nature, Dane pointed out that the cohort describing patients with "COVID" in the radiology report may not include all patients with COVID-19.

"Additionally," the authors of this AJR article continued, "not all patients with <u>thromboembolism</u> undergo confirmative imaging, and some may not experience symptoms. Consequently, the prevalence is likely larger than reported."





B, 46-year-old man who was found to have COVID-19 after presenting with fever, shortness of breath, epigastric pain, and weakness. Axial IV contrastenhanced abdominopelvic CT image shows small left renal infarct (arrow) and nonocclusive aortic, celiac artery, and superior mesenteric artery thrombus (circle). Complete common hepatic artery thrombus (not shown) was also seen. Splenic and renal infarctions (not shown) were present, despite patent visualized associated arteries and veins. Concurrently performed IV contrast-enhanced chest CT angiogram (not shown) showed lobar and segmental pulmonary emboli; probable left lower lobe pulmonary infarction rather than pneumonia; and bilateral, multilobar, patchy but confluent ground-glass opacities typical of COVID-19. Credit: American Roentgen Ray Society (ARRS), American Journal of Roentgenology (AJR)



**More information:** Bari Dane et al, Hypercoagulability in Patients With Coronavirus Disease (COVID-19): Identification of Arterial and Venous Thromboembolism in the Abdomen, Pelvis, and Lower Extremities, *American Journal of Roentgenology* (2020). DOI: 10.2214/AJR.20.23617

## Provided by American Roentgen Ray Society

Citation: Hypercoagulability in patients with COVID-19 (2020, September 24) retrieved 23 May 2024 from <u>https://medicalxpress.com/news/2020-09-hypercoagulability-patients-covid-.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.