

## Bat tick found for the first time in New Jersey

September 9 2020, by Todd Bates



Live larval bat ticks (Carios kelleyi) removed in 2019 from big brown bats in Mercer County, New Jersey. Credit: J. Occi/Rutgers Center for Vector Biology

A tick species associated with bats has been reported for the first time in New Jersey and could pose health risks to people, pets and livestock,



according to a Rutgers-led study in the Journal of Medical Entomology.

This species (Carios kelleyi) is a "soft" <u>tick</u>. Deer ticks, which carry Lyme disease, are an example of "hard" ticks.

"All ticks feed on blood and may transmit pathogens (disease-causing microbes) during feeding," said lead author James L. Occi, a doctoral student in the Rutgers Center for Vector Biology at Rutgers University-New Brunswick. "We need to be aware that if you remove <u>bats</u> from your belfry, attic or elsewhere indoors, ticks that fed on those bats may stay behind and come looking for a new source of blood. There are records of C. kelleyi biting humans."

This soft tick species, a parasite of bats, is known to be in 29 of the 48 contiguous U.S. states, and was confirmed in New Jersey as larvae collected from big brown bats (Eptesicus fuscus) in Mercer and Sussex counties. This is a new addition to the list of New Jersey ticks.

While the public health risk remains unknown, "finding them on New Jersey bats was an unusual event that prompted bat specialists to contact us. Maybe these ticks are becoming more common," said senior author Dina M. Fonseca, a professor and director of the Center for Vector Biology in the Department of Entomology in the School of Environmental and Biological Sciences.

In other states, C. kelleyi has been found infected with microbes that are harmful to people, pets and livestock. There have been reports of this soft tick feeding on humans, and the bat that hosts them regularly roosts in structures such as attics and barns, underscoring the need to learn more about them, the study says.

"This tick belongs to the family Argasidae, known as 'soft ticks' because their body looks leathery and soft," Fonseca said. That is in contrast to



the "hard ticks" (family Ixodidae) that New Jerseyans are more familiar with.

Scientists in the Endangered and Nongame Species Program of the Division of Fish and Wildlife in the New Jersey Department of Environmental Protection found the tick larvae on bats last year. Technically, this is not the first time a soft tick has been reported in New Jersey. In 2001, a related <u>tick species</u>—Carios jersey—was found in amber in Middlesex County. That specimen was 90 million to 94 million years old.

"The next steps are to collect more soft tick specimens and test them for disease-causing microbes," Occi said.

**More information:** James L Occi et al, First Record of Carios kelleyi (Acari: Ixodida: Argasidae) in New Jersey, United States and Implications for Public Health, *Journal of Medical Entomology* (2020). DOI: 10.1093/jme/tjaa189

Provided by Rutgers University

Citation: Bat tick found for the first time in New Jersey (2020, September 9) retrieved 7 May 2024 from <u>https://medicalxpress.com/news/2020-09-jersey-1.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.