

Older women with type 2 diabetes have different patterns of blood use in their brains

September 8 2020, by Laurie Fickman



Stacey Gorniak, associate professor in the University of Houston Department of Health and Human Performance and director of the Center for Neuromotor and Biomechanics Research, is reporting that brains of older women with Type 2 diabetes do not use as much oxygenated blood as those who don't have the disease. Credit: University of Houston



A University of Houston researcher is reporting that the brains of older women with Type 2 diabetes do not use as much oxygenated blood as those who don't have the disease. The research is the first to point to changes in blood use in the brain as the primary reason for diabetesrelated deficits in motor function. It also furthers the understanding of sensory and motor symptoms as a precursor to developing dementia and Alzheimer's diseases, both of which are linked to diabetes.

"It's a pretty significant finding. Typically, when someone presents with a sensory or motor issue along with Type 2 <u>diabetes</u> mellitus, the assumption is that it's the result of peripheral nerve damage in the hands and feet," said Stacey Gorniak, associate professor in the UH Department of Health and Human Performance and director of the Center for Neuromotor and Biomechanics Research. Gorniak published her findings in the journal *Neurophotonics*.

Until now there has been no assumption that something is going on with respect to brain function that is affecting sensory and motor functions in persons living with Type 2 diabetes.

"Emerging evidence has suggested that factors outside of nerve damage due to Type 2 <u>diabetes mellitus</u>, such as impaired cortical blood use, contribute significantly to both sensory and motor deficits in people with diabetes," reports Gorniak.

Nearly 24% of the 40 million people in the United States over the age of 60 live with Type 2 diabetes. Problems with hands, fingers and feet are common side effects of the <u>disease</u> and can lead to a loss of independent living and decline in quality of life.

Gorniak's testing method is unique. Rather than using a typical MRI to monitor the use of oxygenated blood, she opted to use a technique called functional near <u>infrared spectroscopy</u> (fNIRS). The fNIRS is a method



that delivers infrared light into the scalp to measure use of both oxygenated and unoxygenated blood use by the brain. This technique differs from MRI as MRI cannot measure oxygenated <u>blood</u> use. The fNIRS method can be used on persons who cannot have an MRI.

She tested a group of 42 post-menopausal women, over 60, half of whom had Type 2 diabetes, and asked them to perform various exercises with their hands. She chose this group because they are generally at the highest risk for diabetes, heart disease and dementia.

"Our work demonstrates that motor changes in people with diabetes occur independent of <u>sensory impairment</u> and that these changes are unrelated to disease duration and severity. Our data point towards other factors such as changes in muscle and reduced function of the cortex as underlying mechanisms for problems in sensory and motor functions," Gorniak reports.

Her findings, she said, opens research possibilities for other groups of people with the disease, in hopes of finding a way to therapeutically avoid the negative health effects of diabetes.

"We need to see what this looks like in a larger population, including men, and then we can start developing treatments or different ways we could potentially stop these negative impacts of Type 2 diabetes," said Gorniak.

Provided by University of Houston

Citation: Older women with type 2 diabetes have different patterns of blood use in their brains (2020, September 8) retrieved 10 May 2024 from <u>https://medicalxpress.com/news/2020-09-older-women-diabetes-patterns-blood.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.