

Pacemaker need higher in patients with type 2 diabetes

September 16 2020



(HealthDay)—The need for pacemakers is higher among patients with

type 2 diabetes than the general population, according to a study published online Sept. 4 in *Diabetes Care*.

Elina Rautio, from the Solna Karolinska Institutet in Stockholm, and colleagues used data from the Swedish National Diabetes Registry to identify 416,247 patients with type 2 diabetes plus 2,081,235 age- and sex-matched controls selected from the general population. The need for pacemaker implementation was compared between the two groups.

The researchers found that type 2 diabetes was associated with an increased need for pacemaker treatment (hazard ratio, 1.65), which remained significant (hazard ratio, 1.56) after adjusting for age, sex, educational level, marital status, country of birth, and coronary heart disease. Increasing age, hemoglobin A1c, body mass index, diabetes duration, and lipid- and blood pressure-lowering medication were [risk factors](#) for receiving a pacemaker.

"The [medical community](#) should therefore recognize that patients with type 2 diabetes comprise a high-risk group for developing bradyarrhythmias with a subsequent need for pacemaker treatment," the authors write.

More information: [Abstract/Full Text \(subscription or payment may be required\)](#)

Copyright © 2020 [HealthDay](#). All rights reserved.

Citation: Pacemaker need higher in patients with type 2 diabetes (2020, September 16) retrieved 3 May 2024 from <https://medicalxpress.com/news/2020-09-pacemaker-higher-patients-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.