

Patients showing more DNA methylation at some genome sites found to be more likely to fail to respond to metformin

September 17 2020, by Bob Yirka



Credit: CC0 Public Domain

A team of researchers affiliated with multiple institutions in Europe has found a connection between patients who show more DNA methylation

at some genome sites and a higher likelihood of failing to respond to metformin. In their paper published in the journal *Science Translational Medicine*, the group describes their genomic study of hundreds of type 2 diabetes patients and what they learned from them.

Type 2 diabetes is a condition in which a person's body becomes resistant to insulin. As a result, cells throughout the body become unable to absorb glucose, leading to higher blood sugar levels. One of the main treatments for [patients](#) with type 2 diabetes is the drug metformin. It works by making tissues in the body less sensitive to insulin, lowering blood sugar levels. Unfortunately, the drug does not work the way it is supposed to in approximately 30 percent of patients, and some patients experience side effects so severe they are forced to discontinue use of the drug.

In this new effort, the researchers sought to determine if DNA methylation might provide hints of problems with metformin. Methylation is a chemical reaction in cells whereby [methyl groups](#) are added to molecules. The process is important because it can change the activity of DNA segments without changing their sequence. To find out if methylation might be used as a marker to help spot patients who may not respond in positive ways to metformin, they analyzed the genomic data of 345 diabetes type 2 patients.

The work involved parsing epigenetic data from the genomes, which allowed them to see the degree of DNA methylation at different genome sites. They found a connection between more DNA methylation at 11 sites in the genome and a 2.5% increased likelihood of failing to respond to metformin. They also found four other high-level methylation sites that they linked to tripling the likelihood of drug intolerance.

The researchers suggest their results were strong enough to warrant further investigation of the possibility of using methylation testing

before prescribing metformin to [diabetes patients](#).

More information: Sonia García-Calzón et al. Epigenetic markers associated with metformin response and intolerance in drug-naïve patients with type 2 diabetes, *Science Translational Medicine* (2020). [DOI: 10.1126/scitranslmed.aaz1803](https://doi.org/10.1126/scitranslmed.aaz1803)

[Press release](#)

© 2020 Science X Network

Citation: Patients showing more DNA methylation at some genome sites found to be more likely to fail to respond to metformin (2020, September 17) retrieved 21 May 2024 from <https://medicalxpress.com/news/2020-09-patients-dna-methylation-genome-sites.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--