

Many practitioners are not prescribing HIV prevention medication, study finds

September 17 2020



Credit: CC0 Public Domain

Only about 54% of medical practitioners surveyed say they have prescribed pre-exposure prophylaxis, or PrEP, to HIV-vulnerable patients, according to a new study by a Vanderbilt University Medical

Center investigator.

The study, led by Ashley Leech, Ph.D., assistant professor of Health Policy, surveyed 519 practitioners in five major U.S. cities who attended a continuing medical education course on HIV between March and May 2015. The study was published in *PLOS ONE*.

PrEP is a once-daily medication for people without HIV that is a widely accepted method of preventing HIV transmission, but less than 10% of the more than 1 million people vulnerable to HIV are taking PrEP, according to the study.

"With a U.S. policy goal to eradicate HIV by 2030, practitioners are central to ensuring the delivery of PrEP across care settings," Leech said. "Our findings, however, indicate that even among a subset of HIV-focused practitioners, PrEP prescribing is not routine."

The study found that internal medicine prescribers were 1.6 times more likely to prescribe PrEP than infectious disease practitioners, which researchers said was an indicator of how important internal and family medicine practitioners are in assessing and mitigating risk in their patients.

Longstanding confusion or disagreement between HIV and primary care practitioners over who should be responsible for prescribing this preventive medication may partly explain the slow adoption of PrEP among practitioners, Leech said.

Age, years of training, and sex of the independent prescribers were also significantly associated with prescribing experience.

The study also found that practitioners might be more willing to prescribe PrEP as a first clinical step for persons who inject drugs than

previously indicated, with more work needed to effectively communicate with priority groups vulnerable to HIV and to reach individuals who could most benefit from PrEP, according to Leech.

"While a number of factors could impact PrEP prescribing, including patients' low familiarity with the drug or practitioners' lack of opportunity in offering PrEP, in order for the drug to be effective at eliminating HIV in the United States in the next 10 years, the proportion of prescribing needs to increase, with more effort placed on identifying risk across populations and clinical specialties."

More information: Ashley A. Leech et al, Healthcare practitioner experiences and willingness to prescribe pre-exposure prophylaxis in the US, *PLOS ONE* (2020). [DOI: 10.1371/journal.pone.0238375](https://doi.org/10.1371/journal.pone.0238375)

Provided by Vanderbilt University Medical Center

Citation: Many practitioners are not prescribing HIV prevention medication, study finds (2020, September 17) retrieved 12 May 2024 from <https://medicalxpress.com/news/2020-09-practitioners-hiv-medication.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--