


# The distribution range of Ebola virus carriers in Africa may be larger than previously assumed

September 7 2020



Modelled habitat (in red) versus current distribution according to IUCN (schraffiert) of the nine potential reservoir hosts of Zaire ebolavirus a) *Epomops franqueti*, b) *Hypsignathus monstrosus*, c) *Myonycteris torquata*, d) *Eidolon helvum*, e) *Epomophorus gambianus*, f) *Lissonycteris angolensis*, g) *Micropteropus pusillus*, h) *Mops condylurus*, i) *Rousettus aegyptiacus*. Credit: Koch, L.K. et al.

The Ebola virus is considered one of the world's most dangerous pathogens. During the most severe outbreak to date in West Africa, over 11,000 deaths were documented between 2014 and 2016. Single cases are repeatedly reported from Europe as well, which are connected to

previous travels to affected regions. An important source of infection are so-called reservoir hosts that carry the virus without being affected by it. For the various types of the Ebola virus, the most likely involve reservoir hosts are various species of bats and fruit bats.

For the first time, scientists investigated where nine of such bat and fruit bat species may encounter [suitable habitats](#) and climatic conditions in Africa. "Zaire ebolavirus is one of the most dangerous Ebola viruses. It kills up to 88 percent of those infected with it. To prevent or curb outbreaks of this [virus](#), it is essential to know exactly where potential hotspots of infection may lurk," explains parasitologist Prof. Dr. Sven Klimpel of the Goethe University in Frankfurt and the Senckenberg Biodiversity and Climate Research Center.

Based on ecological niche modeling, his team was able to show that the respective bat and fruit bat species are able to thrive in West and East Africa, including large parts of Central Africa. A wide belt of potential habitats extends from Guinea, Sierra Leone, and Liberia in the west across the Central African Republic, the Republic of the Congo and the Democratic Republic of the Congo to Sudan and Uganda in the East. A few of the studied bats and fruit bats may even occur in the eastern part of South Africa.

In a second step, the researchers compared the potential habitats with range maps of the bat and fruit bat species that were generated by the International Union for the Conservation of Nature (IUCN) on the basis of observations of these animals. In addition, the team considered where a Zaire ebolavirus endemic has broken out in the past. The results were surprising: "The modeled habitats of the Zaire ebolavirus hosts are larger than their previously known ranges. It is possible that the bats and [fruit bats](#) have not yet been able to reach habitats beyond these ranges due to the presence of certain barriers," says Klimpel.

"Another, more worrying explanation could be that science has hitherto underestimated the range of Ebola-transmitting bat and fruit bat species. In this case, the models would provide a more realistic picture," explains Dr. Lisa Koch, the study's lead author from Goethe University. Regions affected by Ebola outbreaks often suffer not only from health effects, but also from economic and social effects of the epidemic. The study's findings suggest to keep a closer eye on diseases that occur in the modeled ranges of the reservoir hosts and to inform the public about potential Ebola infections, ultimately alleviating the consequences of an epidemic.

With regard to Europe, Klimpel states: "Ebola viruses, just like the SARS-CoV-2 (Coronavirus), are viruses from the animal kingdom that can be transmitted to humans. It can be expected that diseases of this type will occur more frequently in the future, since humans have increasing contact with wild animals, and globalization facilitates the spread of viruses around the world. In Europe, with its overall efficient health system, Ebola infections are certainly going to remain isolated incidents in the future. Nonetheless, in view of these trends it would be beneficial to intensively train and further educate physicians and nursing personnel in the treatment of tropical infectious diseases in our latitudes as well."

**More information:** Lisa K. Koch et al. Bats as putative Zaire ebolavirus reservoir hosts and their habitat suitability in Africa, *Scientific Reports* (2020). [DOI: 10.1038/s41598-020-71226-0](https://doi.org/10.1038/s41598-020-71226-0)

Provided by Senckenberg Research Institute and Natural History Museum

Citation: The distribution range of Ebola virus carriers in Africa may be larger than previously

assumed (2020, September 7) retrieved 30 April 2024 from  
<https://medicalxpress.com/news/2020-09-range-ebola-virus-carriers-africa.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.