

Science of champion runners: Inside the body of elite endurance athletes

September 30 2020, by Andy Galbraith



Credit: AI-generated image ([disclaimer](#))

The 40th anniversary of the London Marathon takes place on Sunday, October 4 2020. Athletes will run on a closed-loop circuit around St James's Park before finishing on The Mall. This year's lineup includes current champions Eliud Kipchoge and Brigid Kosgei. These athletes can run for more than two hours at speeds an average person could maintain

for only a matter of seconds. So what makes them so fast?

Years of training have brought about bodily adaptations that enable elite endurance athletes to perform at the top of their game. Let's first look at the all-important cardiovascular system, which is adapted to enable a regular supply of oxygenated blood to the working muscles.

Strong heart

Regular training results in an increase in the [thickness](#) of the muscle forming the [heart](#) wall, in particular, the left side of the heart. This is the side of the heart responsible for circulating oxygenated blood around the body and to the muscles during [exercise](#).

Increased thickness of the left ventricle wall (the lower chamber of the heart), enables endurance athletes to pump more blood out of their heart each time it beats. We call this the "stroke volume."

A regular adult, exercising at their maximum level, can have a stroke volume of [120ml](#). However, because of the increased thickness of the heart wall, endurance athletes have a stroke volume during maximal exercise of around [200ml](#).

An endurance athlete's heart can beat 200 times every minute during maximal exercise, resulting in these athletes being able to circulate close to [40 liters](#) of blood around the body every minute. The amount of blood the heart pumps in one minute is called cardiac output. In elite endurance runners, it can be nearly twice that of an average adult during maximal exercise.

VO2 max

It is not just a strong heart that is key to these athletes' success. Getting enough oxygen to the working muscles is also vital.

Activities lasting longer than a couple of minutes mainly rely on [aerobic metabolism](#) (the process by which we use oxygen to turn fuel, such as fats and sugars, into energy), making a constant supply of oxygen crucial to success. Elite marathon runners can breathe around [200 liters](#) of air per minute during maximal exercise. This combined ability of the lungs and heart to take in and transport oxygen means that [elite athletes](#) have a very high VO₂ max—the maximum amount of oxygen the body can use in a minute. VO₂ max is regarded as one of the key determinants of [endurance performance success](#).

VO₂ max values in an average adult are around [30-45](#) milliliters of oxygen per kilogram of body weight per minute. But in elite endurance athletes, VO₂ max increases to [65-80 ml/kg/min](#).

Running economy

It's not only the size of an athlete's engine (VO₂ max) that's important for success, just like a car, the economy of the engine also plays an important role. Most marathons are run at around [75-85% of VO₂ max](#), meaning the ability to work efficiently at lower exercise intensities is also key to success. In running, this is measured by "running economy."

Athletes with good running economy need less oxygen to run at a given speed than their competitors, preserving vital energy for later in the race. Elite endurance runners display incredibly low running economy values, showing their ability to move at fast speeds while using a much lower amount of oxygen than the average person.

At a given speed, an average person may typically need 220ml of oxygen per kilogram of body weight to run one kilometer. Elite marathon

runners are more economical, perhaps requiring as little as [180ml](#) of [oxygen](#) per kilogram of body weight to cover the same distance.

Lactate threshold

An important factor in running a fast marathon race is the ability to run at the fastest possible speed without becoming tired. This optimal speed or "threshold" is related to several changes that happen in our body as we exercise, including the build-up of chemicals in the blood. Blood lactate is one such substance that accumulates in the blood during exercise.

[Lactate threshold](#) is a term used in exercise physiology to describe specific changes (or breakpoints) to this substance during exercise.

Blood lactate is often incorrectly thought of as a waste product and mistakenly thought to be responsible for muscle fatigue, but it is now recognized as an [important energy source](#). Still, exercise above the lactate threshold is associated with more rapid fatigue and, as such, the speed at lactate threshold is a [powerful predictor](#) of endurance performance.

A higher lactate threshold will enable a higher [running speed](#) to be sustained without [blood](#) lactate accumulating, enabling the running [speed](#) to be maintained for an extended period. Elite male and female marathon runners have reported lactate threshold speeds of [18-21km/h](#)

As the runners lap St James's Park on Sunday, all the above factors will contribute to their success—they will be the [key determinants](#) of the runners' performance.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Science of champion runners: Inside the body of elite endurance athletes (2020, September 30) retrieved 27 April 2024 from <https://medicalxpress.com/news/2020-09-science-champion-runners-body-elite.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.