


Researchers study how tumor cells divide in the crowd

September 2 2020



A mini-tumor of human breast epithelial cells (MCF-7). A dividing cell indicated in green. Credit: Dr. Elisabeth Fischer-Friedrich

Scientists led by Dr. Elisabeth Fischer-Friedrich, group leader at the Excellence Cluster Physics of Life (PoL) and the Biotechnology Center TU Dresden (BIOTEC) studied how cancer cells are able to divide in a crowded tumor tissue and connected it to the hallmark of cancer progression and metastasis, the epithelial-mesenchymal transition (EMT).

Most [animal cells](#) need to become spherical in order to divide. To achieve this round shape, the [cells](#) must round up and deform their neighboring cells. In a growing [tumor](#) tissue, the tumor cells need to divide in an environment that is becoming more crowded than the healthy tissue. This means that the dividing tumor cells likely need to generate much higher mechanical forces to round up in such a densely packed surrounding. Yet, tumor cells seem to be adapted to overcome these difficulties. Scientists led by Dr. Elisabeth Fischer-Friedrich were curious how do the tumor cells gain this enhanced ability to deal with the crowded tumor environment?

The researchers found that the EMT could be one of the answers. What is it exactly? "EMT or [epithelial-mesenchymal transition](#) is a hallmark of cancer progression," says Kamran Hosseini, Ph.D. student who performed the experiments. It is a cell transformation during which tumor cells lose their asymmetric organization and detach from their neighbors, gaining the ability to migrate into other tissues. This, together with other factors, allows tumors to metastasize, i.e., move into the blood and lymphatic vessels and ultimately colonize other organs.

"So far, EMT has been mainly linked to this enhanced cell dissociation and cell migration. Our results suggest that EMT might also influence [cancer cells](#) by promoting successful rounding and cell division. These results point towards a completely new direction of how EMT could promote metastasis of carcinoma in the body," explains Kamran Hosseini.

Just as we test the ripeness of the fruits by squeezing them gently with our hands, the scientists examined the mechanical properties of individual human cells. Except, they squished the cells using an atomic force microscope. This state-of-the-art setup measured properties such as cell stiffness and cell surface tension before and after the EMT. In addition, the group of Dr. Elisabeth Fischer-Friedrich in collaboration with Dr. Anna Taubenberger (BIOTEC, TU Dresden) and Prof. Carsten Werner (IPF, Dresden) cultured mini-tumors and trapped them inside elastic hydrogels to check how mechanical confinement affects cell rounding and division of tumor cells.

The authors identified changes in rounding and growth of the tumor. EMT influenced the cancer cells in two contrasting ways. The dividing [tumor cells](#) became stiffer while surrounding non-dividing cells became softer. Furthermore, the researchers found hints that the observed mechanical changes could be linked to the increased activity of a protein called Rac1, a known regulator of the cytoskeleton.

"Our findings will not only provide important results to the field of cell biology but may also identify new targets for [cancer](#) therapeutics," says Dr. Elisabeth Fischer-Friedrich.

More information: Kamran Hosseini et al. EMT-Induced Cell-Mechanical Changes Enhance Mitotic Rounding Strength, *Advanced Science* (2020). [DOI: 10.1002/advs.202001276](https://doi.org/10.1002/advs.202001276)

Provided by Dresden University of Technology

Citation: Researchers study how tumor cells divide in the crowd (2020, September 2) retrieved 25 April 2024 from <https://medicalxpress.com/news/2020-09-tumor-cells-crowd.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.