


New research finds low bone health testing rates after prostate cancer treatment

October 7 2020

JOURNAL OF THE NATIONAL COMPREHENSIVE CANCER NETWORK



NCCN Guidelines®

• CHRONIC MYELOID LEUKEMIA

NCCN Guidelines® Insights

• COLORECTAL CANCER SCREENING

ONCOLOGY WATCH

Flirting With Disaster: COVID-19 and Cancer
Tempero M

COMMENTARY

The Home is the New Cancer Center
Handley NR, Bekelman JE, and Binder AF

THE LAST WORD

Time for More Inclusive Cancer Trials
Dirix P, Wyld L, Paluch-Shimon S, and Poortmans P

MOLECULAR INSIGHTS IN PATIENT CARE

A Cryptic *BCR-PDGFRB* Fusion Resulting in a Chronic Myeloid Neoplasm With Monocytosis and Eosinophilia: A Novel Finding With Treatment Implications
Gupta SK, Jain N, Tang G, et al

REVIEW

Treatment Outcomes in Chronic Myeloid Leukemia: Does One Size Fit All?
Garcia-Horton A and Lipton JH

SPECIAL FEATURE

Available at JNCCN.org

Considerations for Use of Hematopoietic Growth Factors in Patients With Cancer Related to the COVID-19 Pandemic
Griffiths EA, Alwan LM, Bachiashvili K, et al

ORIGINAL RESEARCH

Performance Status Restriction in Phase III Cancer Clinical Trials
Abi Jaoude J, Kouzy R, Mainwaring W, et al

Analysis of Sentinel Node Biopsy and Clinicopathologic Features as Prognostic Factors in Patients With Atypical Melanocytic Tumors
Maurichi A, Miceli R, Patuzzo R, et al

Small Renal Masses With Tumor Size 0 to 2 cm: A SEER-Based Study and Validation of NCCN Guidelines
Pecoraro A, Rosiello G, Luzzago S, et al

Association Between Clinical Value and Financial Cost of Cancer Treatments: A Cross-Sectional Analysis
Mitchell AP, Tabatabai SM, Dey P, et al

Relationship Between Quality of Life and Survival in Patients With Pancreatic and Periampullary Cancer: A Multicenter Cohort Analysis
Mackay TM, Latenstein AEJ, Sprangers MAG, et al

A Pilot Study of a Comprehensive Financial Navigation Program in Patients With Cancer and Caregivers
Watabayashi K, Steelquist J, Overstreet KA, et al

Contemporary Population-Based Analysis of Bone Mineral Density Testing in Men Initiating Androgen Deprivation Therapy for Prostate Cancer
Hu J, Aprikian AG, Vanhuysse M, and Dragomir A

New research in the October 2020 issue of *JNCCN—Journal of the National Comprehensive Cancer Network* finds the rate of bone mineral density (BMD) testing in people with prostate cancer undergoing androgen deprivation therapy (ADT) has improved in recent years, but remains low. ADT is considered a cornerstone of treatment for high-risk or advanced prostate cancer and is used in nearly half of all prostate cancer patients. However, it can result in preventable side effects like osteoporosis and bone fractures. Despite clinical recommendations that call for BMD testing in ADT recipients, only 23.4% of the patients studied received testing in 2015. That is up from just 4.1% in 2000.

"Although we expected BMD testing rates to be fairly low given the prior literature, we were somewhat surprised that they didn't go up more in recent years," said senior author Alice Dragomir, MSc, Ph.D., McGill University in Montreal, Quebec who worked with Armen G. Aprikian, MD, Marie Vanhuyse, MD, MSc, and Jason Hu, MSc, also from McGill. "Bone density testing helps doctors evaluate fracture risk and identify which patients would benefit from additional monitoring and interventions like lifestyle changes and/or medications. Perhaps the low rate of testing will change in the coming years thanks to renewed attention on bone health issues in the clinical oncology community. It may be interesting to re-examine BMD testing rates in a few years."

The researchers used the Régie de l'assurance maladie du Québec (RAMQ)—a Canadian public healthcare administrative database—to review patient demographic and billing information for 22,033 people with [prostate cancer](#) who began receiving ADT between January 2000 and December 2015. Of those, 3,910 (17.8%) received a BMD [test](#) at any point during the study period. The largest increase in testing rates

occurred around 2003 and 2004, coinciding with the publication of several articles and guidelines recommending BMD screening in this set of patients. People age 80 and older, with metastatic disease, or living in rural areas were less likely to be screened.

"While we have known for many years that the androgen deprivation therapies used to treat [prostate cancer](#) carry an increased risk of osteoporosis, this study identifies specific populations that might not undergo recommended screening prior to hormone-based therapies," commented Joshua M. Lang, MD MS, Associate Professor of Medicine, Carbone Cancer Center, University of Wisconsin. "These populations are especially vulnerable, including our older patients located in rural areas of the country."

Dr. Lang, a Member NCCN Guidelines Panel for Prostate Cancer, who was not involved in this research, continued: "The importance of screening is even more critical given the availability of medications that can slow or reverse osteoporosis. The NCCN Guidelines for Prostate Cancer specifically recommend screening for these patients and this report demonstrates that more work is needed to advocate for and implement screening of vulnerable patient populations."

More information: Jason Hu et al, Contemporary Population-Based Analysis of Bone Mineral Density Testing in Men Initiating Androgen Deprivation Therapy for Prostate Cancer, *Journal of the National Comprehensive Cancer Network* (2020). [DOI: 10.6004/jnccn.2020.7576](https://doi.org/10.6004/jnccn.2020.7576)

Provided by National Comprehensive Cancer Network

Citation: New research finds low bone health testing rates after prostate cancer treatment (2020, October 7) retrieved 26 April 2024 from <https://medicalxpress.com/news/2020-10-bone-health->

[prostate-cancer-treatment.html](#)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.