

## What do breast cancer cells feel inside the tumour?

October 27 2020



Cancer cell during cell division. Credit: National Institutes of Health

Using a new technique, a team of McGill University researchers has found tiny and previously undetectable 'hot spots' of extremely high stiffness inside aggressive and invasive breast cancer tumors. Their



findings suggest, for the first time, that only very tiny regions of a tumor need to stiffen for metastasis to take place. Though still in its infancy, the researchers believe that their technique may prove useful in detecting and mapping the progression of aggressive cancers.

"We are now able to see these features because our approach allows us to take measurements within living, intact, 3-D tissues," says Chris Moraes, from McGill University's Department of Chemical Engineering, a Canada Research Chair and senior author on a recent research paper in *Nature Communications*. "When <u>tissue samples</u> are disrupted in any way, as is normally required with standard techniques, signs of these 'hot spots' are eliminated."

## "Smart" hydrogels provide information about cancer progression

The researchers built tiny hydrogel sensors that can expand on demand, much like inflating balloons the size of individual cells, and placed them inside 3-D cultures and mouse models of breast <u>cancer</u>. When triggered, the expansion of the hydrogel can be used to measure very local stiffness inside the tumor.

This unusual technique, developed through a collaboration between McGill's Department of Chemical Engineering and the Rosalind and Morris Goodman Cancer Research Centre at McGill, allows the researchers to sense, from the perspective of a cancer cell, what is going on in their surrounding environment.

## What cells sense drives their behavior

"Human cells are not static. They grab and pull on the tissue around them, checking out how rigid or soft their surroundings are. What cells



feel around them typically drives their behavior: immune cells can activate, stem cells can become specialized, and cancer cells can become dangerously aggressive," explains Moraes. "Breast cancer cells usually feel surroundings that are quite soft. However, we found that cancer <u>cells</u> inside aggressive tumors experienced much harder surroundings than previously expected, as hard as really old and dried up gummy bears."

The researchers believe that their findings suggest new ways in which cell mechanics, even at the early stages of breast cancer, might affect disease progression.

"Developing methods to analyze the mechanical profiles in 3-D tissues may better predict patient risk and outcome," says Stephanie Mok, the first author on the paper and a Ph.D. candidate in the Department of Chemical Engineering. "Whether these 'hot spots' of stiffness are really causing cancer progression rather than simply being correlated with it remains an open, but critically important question to resolve."

**More information:** Stephanie Mok et al, Mapping cellular-scale internal mechanics in 3D tissues with thermally responsive hydrogel probes, *Nature Communications* (2020). DOI: 10.1038/s41467-020-18469-7

Provided by McGill University

Citation: What do breast cancer cells feel inside the tumour? (2020, October 27) retrieved 4 May 2024 from <u>https://medicalxpress.com/news/2020-10-breast-cancer-cells-tumour.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is



provided for information purposes only.