

# How to prevent COVID-19 'superspreader' events indoors this winter

October 28 2020, by Gabriel Wainer

---



Credit: AI-generated image ([disclaimer](#))

The second wave of COVID-19 has arrived. [Cases are spiking in most European countries](#), Canada is seeing the number of daily cases [exceed the first peak in April/May](#) and the United States is facing a [potential third wave](#).

Some countries like Brazil, India and Argentina are still in the [first wave](#). Others like New Zealand, Vietnam and Singapore have [contained the virus](#). But the second wave has arrived in the northern hemisphere, as we confront the upcoming winter.

In most countries, the number of deaths is [lower than in the first wave](#); we have learned more about the virus and know that the second wave is driven by younger people, which have a lower death rate. Nevertheless, the long-term effects of this wave are yet to be seen.

Mathematical modeling is used to predict the spread of diseases like COVID-19. These models must now take into consideration people's changing behaviors and the approaching winter.

## **Unknown long-term effects**

There is evidence that COVID-19 infection can lead to [serious heart problems](#), which are currently [being studied](#). Some people also experience damage to the [kidneys, lungs and brain](#). There have also been reports of [sudden deafness](#) and "[brain fog](#)."

We still do not know if [there is immunity after infection](#) or whether [the disease will become endemic](#).

The second wave might have been triggered by [lockdown fatigue and feelings of invincibility of young population](#). But [coping mechanisms like denial and rationalism](#), combined with conflicting public health messages, can lead to [population disobedience](#) and trigger a large number of cases.

## **Superspreader events**

We have learned important aspects about the spread of COVID-19. The

pandemic has largely been driven by [superspreader events](#), in which a single individual directly infects several contacts.

The reproductive number—known as  $R_0$ —of SARS-CoV-2 is between two and three, meaning that, on average, an individual infects between two and three individuals. The variation in the distribution—known as  $k$ —has showed that some individuals generate [a large number of secondary cases](#): less than [20 percent of individuals produce over 80 percent of infections](#).

There have been numerous examples of superspreader events such as the [outbreak associated with air conditioning in a restaurant in Guangzhou, China](#), the exposure of individuals at a [choir practice in Skagit County, Wash.](#), [the recent Rose Garden ceremony at the White House](#) and a [spinning class in Hamilton, Ont.](#)

## The role of aerosols

We also know that the disease is driven by aerosols, which are [small particles](#) made of [saliva or respiratory fluid whose size is a few microns](#). The U.S. Centers for Disease Control and Prevention states that the disease mainly spreads through [close contact](#). They also state that touching contaminated surfaces (known as fomites) is not thought to be a common way that COVID-19 spreads.

Since April, we've learned that [aerosols and indoor transmission are the main sources of infection](#). There is evidence that the virus can remain [in the air from 15 minutes up to three hours](#) and transmission indoors may be around [20 times higher than outdoors](#).

Numerous studies show that transmission is due to indoor spreading via aerosols. A recent study of [1,200 outbreaks](#), shows that with the exception of two incidents, most outbreaks were indoor and driven by


homes and family contacts.

## **Paying attention to ventilation**

To help reduce the probability of superspreading events indoors, [we need to pay attention to ventilation](#). Ideally, we should conduct most activities outdoors, but when not possible, air quality should be guaranteed.

Ventilation standards like [ASHRAE and MERV-13 should be mandatory](#), and when these are not possible, portable air purifiers with HEPA filters [can reduce the probability of infection](#).

We can also use carbon dioxide detectors to [check the quality of air indoors](#). And with the cold weather approaching, controlling [the level of carbon dioxide](#) and humidity indoors can reduce the probability of contagion.



Modelling results that show that intermittent lockdowns flatten the curve, but individual disobedience affects the spread rate of the disease. Credit: Román Cárdenas, Gabriel Wainer, Author provided

## Simulating behaviors

My research team works on flexible modeling methodologies that focus on [spatial models](#) and can easily be adapted to include the most recent research results.

We built a variety of models to study various aspects of the spread of SARS-CoV-2, including connectivity or mobility factors, mortality rates and interventions. We can study the potential outcomes of complex

government policies, such as enforcing the use of masks or physical distancing.

We can also explore mechanisms to combine the base epidemiological models with information about [the surrounding environment \(buildings, transportation, open spaces\)](#). The influence of the spatial location on the model dynamics can be easily visualized to facilitate more efficient decision-making.

## Simulating environments

A second aspect of our research investigates how to integrate advanced 3-D building models with computer models of how the disease spreads indoors. The research focuses on the spread of viral particles through heating, ventilation and cooling (HVAC) systems in buildings. By studying the effect of [humidity levels indoors](#) or the effects of viral particle accumulation in rooms, we can better quantify the risk of COVID-19 and the effectiveness of measures to mitigate the risk.

Our advanced models use carbon dioxide as a proxy for the disease, because the sensors are cheap and sensitive to HVAC systems and room dimensions. These models can help study changes in ventilation, accumulation of viral particles over time and the influence of [voice tone levels](#) in a room.

These simulations can help to assess the risk of crowded and confined spaces, as well as [close contacts](#).

Our next efforts should include modeling long-term side effects and paying attention to individual interactions to study the effects of lockdowns, economic aspects and uncertainty in human behavior. Advanced tools to [model these social aspects are needed](#) to identify problems and solutions. Social aspects—including [leadership, social](#)

[environments, stress and coping behaviors](#)—should not be neglected in future research efforts.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: How to prevent COVID-19 'superspreader' events indoors this winter (2020, October 28) retrieved 24 April 2024 from <https://medicalxpress.com/news/2020-10-covid-superspreader-events-indoors-winter.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.