

Discovery holds potential for reversing vision loss

October 5 2020



Credit: CC0 Public Domain

An Oklahoma Medical Research Foundation discovery could pave the way for therapies to reverse vision loss common in premature infants and adults.



In a new study appearing in the *Proceedings of the National Academy of Sciences*, OMRF scientists have identified a compound that could give birth to therapies for a host of eye diseases that include <u>retinopathy of prematurity</u> and <u>diabetic retinopathy</u>.

"Potentially, even patients with advanced disease progression could see their fortunes turned around," said Courtney Griffin, Ph.D., the senior author of the study.

Several eye disorders occur when <u>blood vessels</u> grow out of control in the retina, the tissue that lines the back of the eye. In these forms of retinopathy, a web of vessels blocks light from reaching the retina, which is how we see. The overgrowth causes vision issues that can advance to total blindness.

Retinopathy in premature babies—linked to high oxygen levels in NICU incubators that interrupts normal vessel development in the eye—often resolves naturally over time. But not always. In those cases, and in adult diseases like diabetic retinopathy, vision damage can be irreversible.

Griffin and Chris Schafer, Ph.D., study the development of blood vessels. Schafer, a postdoctoral researcher, thought there might be clues to thinning this tangle of vessels if they analyzed a different set of vessels that naturally regress and disappear in mice soon after birth.

When they studied newborn mice, the OMRF researchers found that levels of a specific class of cellular proteins crashed as the mice experienced normal blood <u>vessel</u> loss in the eye.

"Dr. Schafer hypothesized that these cellular proteins might be an important 'off switch' to eliminate these vessels in a neonatal model," said Griffin, who holds the Scott Zarrow Chair in Biomedical Research at OMRF. "This is a new way of approaching these diseases. The current



methods—invasive surgeries or life-long injections into the eye—only prevent the <u>disease</u> from advancing and often have serious complications."

Schafer identified an experimental compound that disables the proteins. It allowed him to flip the switch and test the idea.

"We wanted to trick blood vessels in diseased mice into thinking they were supposed to be regressing and naturally dying off," said Schafer. "This appears to be what happened."

Even more encouraging, Griffin said, the compound only impacted abnormal blood vessels with slow blood flow. The normal vessels needed in a healthy eye were spared.

The findings open the door to tailored therapies to reverse vision loss. It may also have implications in shrinking tumors that contain abnormal <u>blood</u> vessels in other parts of the body.

"We've shown that once these abnormal vessels have formed in the young eye, they're susceptible to being treated," said Griffin. The research team will now study the compound in models of adult eye diseases.

"More research is needed, but this could be a major advance in treatment for vision loss in patients of all ages," Griffin said.

More information: Christopher M. Schafer el al., "An inhibitor of endothelial ETS transcription factors promotes physiologic and therapeutic vessel regression," *PNAS* (2020). www.pnas.org/cgi/doi/10.1073/pnas.2015980117



Provided by Oklahoma Medical Research Foundation

Citation: Discovery holds potential for reversing vision loss (2020, October 5) retrieved 3 May 2024 from https://medicalxpress.com/news/2020-10-discovery-potential-reversing-vision-loss.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.