

Lithium decreases the risk of developing severe hyponatremia

October 12 2020, by Cecilia Bodén



Credit: CC0 Public Domain

Researchers at Karolinska Institutet have conducted a study in which lithium treatment was compared between 11,213 individuals hospitalized due to hyponatremia and 44,801 controls. Interestingly, lithium treatment was half as common among individuals with hyponatremia. The results were recently published in *Journal of Psychopharmacology*.

Hyponatremia is a potentially serious disorder that may cause balance disturbances, confusion and even death. The authors have established a cohort that includes all of the 11,000 individuals hospitalized due to

hyponatremia in Sweden over 10 years. The [cohort](#) has been used to study the associations between different groups of drugs and the risk to suffer from hyponatremia that demand inpatient care. Most drugs are associated with an increased risk. However, according to the present study, lithium treatment was associated with a risk that was halved.

"Even though the study confirmed our hypothesis, we were surprised over the marked protective effect associated with lithium treatment," says Henrik Falhammar, lead author of the study and associate professor at the Department of Molecular Medicine and Surgery, Karolinska Institutet.

"The most common reason for development of hyponatremia is fluid retention due to an inadequate release of vasopressin. On the contrary, lithium treatment often leads to the development of nephrogenic diabetes insipidus that result in impairment of the ability to concentrate the urine and large volumes of urine. Therefore, we hypothesized that [lithium treatment](#) might be protective in this regard. Lithium treatment is indeed associated with several side effects. But nevertheless, it is very important for many patients suffering from bipolar disease. Therefore, it was very satisfying to note that [lithium](#) in this regard actually may have a favorable effect," says Buster Mannheimer, associate professor at the Department of Clinical Science an Education and senior researcher for the project.

More information: Henrik Falhammar et al. Reduced risk for hospitalization due to hyponatraemia in lithium treated patients: A Swedish population-based case–control study, *Journal of Psychopharmacology* (2020). [DOI: 10.1177/0269881120937597](https://doi.org/10.1177/0269881120937597)

Provided by Karolinska Institutet

Citation: Lithium decreases the risk of developing severe hyponatremia (2020, October 12)
retrieved 18 April 2024 from
<https://medicalxpress.com/news/2020-10-lithium-decreases-severe-hyponatremia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.