

Nursing home residents with cognitive impairment more likely to be admitted to hospital

October 19 2020



EMERGENCY VISITS FOR NURSING HOME RESIDENTS

MORE LIKELY TO BE DISCHARGED
RESIDENTS WITH:

- FALLS, TRAUMA OR FRACTURE
- BLEEDING
- PAIN

MORE LIKELY TO BE HOSPITALIZED
RESIDENTS WITH:

- COGNITIVE IMPAIRMENT
- RESPIRATORY DIFFICULTIES

 Regenstrief Institute

The infographic features a background image of an elderly man with his hands clasped, looking down thoughtfully. The text is overlaid on a dark grey header and a light blue background.

Transfers from the nursing home to the emergency department (ED) or the hospital can have negative longer-term impact on the health of older adults. A new study from Regenstrief Institute and Indiana University School of Medicine looked at which residents were most likely to be admitted to the hospital after a trip to the ED with the hope of identifying areas to improve care and reduce unnecessary transfers. Credit: Regenstrief Institute

Transfers from the nursing home to the emergency department (ED) or the hospital can have negative longer-term impact on the health of older adults. A new study from Regenstrief Institute and Indiana University School of Medicine looked at which residents were most likely to be admitted to the hospital after a trip to the ED with the hope of identifying areas to improve care and reduce unnecessary transfers.

The study team analyzed hundreds of transfers from the nursing facility to the ED and found the factors most associated with discharge back to the facility were:

- Falls, trauma or fracture
- Bleeding (non-gastrointestinal)
- Pain

Residents who had the following characteristics were more likely to be admitted to the hospital following an ED visit:

- Cognitive impairment
- Respiratory difficulties

"Transfers of care can be very difficult on the patient. Our goal in analyzing this data is to identify new opportunities to improve care, whether it's preventing transfers to the emergency department or finding ways to avoid hospital admissions," said Kathleen Unroe, M.D., MHA, first author of the paper, Regenstrief research scientist and associate professor of medicine at IU School of Medicine. "This information helps us understand what ailments could potentially be treated in the nursing [home](#), avoiding a potentially risky transfer of care."

About one quarter of [nursing facility](#) residents will experience a transfer

to the hospital over the course of a year. There is a significant likelihood of reduced functioning and overall negative impact on their health after discharge from the [hospital](#).

Analyzing the data for hospital admission

The research team looked at 867 long-stay nursing home facility residents enrolled in the OPTIMISTIC project. OPTIMISTIC (Optimizing Patient Transfers, Impacting Medical quality, Improving Symptoms—Transforming Institutional Care) was a demonstration project funded by the Centers for Medicare and Medicaid that successfully reduced potentially avoidable hospitalizations of people in nursing homes.

"Through OPTIMISTIC, we've determined that we can reduce avoidable hospitalizations. The next step is to gain a more nuanced understanding of transfers," said Susan Hickman, Ph.D., senior author on the paper and director of Indiana University Center for Aging Research at Regenstrief Institute. "This helps us find opportunities for improvement from both the facility perspective and the [emergency department](#) perspective."

Researchers say these results suggest one area for exploration may include targeted resources and protocols focused on treatment for falls or symptom management within nursing homes. Another potential area of improvement is the creation of strategies to ensure that providers in the ED have more information when they are assessing the residents, as well as an understanding of the care capabilities of the nursing home to increase the number of discharges back to the facility and avoid hospitalizations.

More information: Kathleen T. Unroe et al, Long-Stay Nursing Facility Resident Transfers: Who Gets Admitted to the Hospital?, *Journal of the American Geriatrics Society* (2020). [DOI](#):

[10.1111/jgs.16633](https://doi.org/10.1111/jgs.16633)

Provided by Regenstrief Institute

Citation: Nursing home residents with cognitive impairment more likely to be admitted to hospital (2020, October 19) retrieved 23 July 2024 from <https://medicalxpress.com/news/2020-10-nursing-home-residents-cognitive-impairment.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.