

Pathogens in the mouth induce oral cancer

October 1 2020



Dental care. Credit: Pixabay user Vanessafrazao

Pathogens found in tissues that surround the teeth contribute to a highly aggressive type of oral cancer, according to a study published 1st October in the open-access journal *PLOS Pathogens* by Yvonne Kapila



of the University of California, San Francisco, and colleagues. In addition, the study showed that oral cancer formation mediated by the pathogens is inhibited by a bacteriocin—an antimicrobial and probiotic peptide that is produced by bacteria.

Head and neck <u>squamous cell carcinoma</u> (HNSCC) is one of the most common cancers worldwide. Oral squamous cell carcinoma (OSCC), a subset of HNSCC, accounts for 90% of all oral malignancies, and it has a poor five-year survival rate that has not changed in decades. Risk factors, including smoking, <u>alcohol consumption</u> and human papilloma virus infection, alone have not been sufficient to explain the incidence and aggressive nature of OSCC. Other factors such as <u>oral pathogens</u> may play an important role in OSCC tumor development, progression and metastasis, yet this has not been well explored. Kapila and her collaborators tested whether OSCC is promoted by periodontal pathogens (i.e., those affecting the structures surrounding and supporting the teeth).

They found that three types of periodontal pathogens (Porphyromonas gingivalis, Treponema denticola, and Fusobacterium nucleatum) enhanced OSCC cell migration, invasion, and tumor formation in mice. These effects were mediated by interactions between two signaling pathways: integrin/FAK and TLR/MyDD88. The pathogen-mediated processes were inhibited by treatment with nisin—a bacteriocin and a commonly used food preservative. According to the authors, this study offers the first direct evidence that a bacteriocin inhibits oral cancer formation mediated by periodontal pathogens. Moreover, the findings suggest that nisin could have broad therapeutic potential as an antimicrobial and anticancer agent, and as an inhibitor of pathogen-mediated cancer formation.

The authors conclude, "Since a probiotic bacteriocin peptide, nisin, rescues this pathogen-mediated carcinogenesis, these findings could



advance treatment for oral cancer and establish a novel paradigm for cancer treatment focused on antimicrobial-based therapies."

More information: Kamarajan P, Ateia I, Shin JM, Fenno JC, Le C, Zhan L, et al. (2020) Periodontal pathogens promote cancer aggressivity via TLR/MyD88 triggered activation of Integrin/FAK signaling that is therapeutically reversible by a probiotic bacteriocin. *PLoS Pathog* 16(10): e1008881. <u>doi.org/10.1371/journal.ppat.1008881</u>

Provided by Public Library of Science

Citation: Pathogens in the mouth induce oral cancer (2020, October 1) retrieved 3 May 2024 from <u>https://medicalxpress.com/news/2020-10-pathogens-mouth-oral-cancer.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.