

Postpartum depression may persist three years after giving birth

October 27 2020



Credit: CC0 Public Domain

A National Institutes of Health study of 5,000 women has found that approximately 1 in 4 experienced high levels of depressive symptoms at some point in the three years after giving birth. The rest of the women

experienced low levels of depression throughout the three-year span. The study was conducted by researchers at NIH's Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD). It appears in the journal *Pediatrics*.

The American Academy of Pediatrics recommends that pediatricians screen mothers for postpartum depression at well-child visits at one, two, four and six months after childbirth. Researchers identified four trajectories of postpartum [depressive symptoms](#) and the factors that may increase a woman's risk for elevated symptoms. The findings suggest that extending screening for postpartum depressive symptoms for at least two years after childbirth may be beneficial, the authors write.

"Our study indicates that six months may not be long enough to gauge depressive symptoms," said Diane Putnick, Ph.D., the primary author and a staff scientist in the NICHD Epidemiology Branch. "These long-term data are key to improving our understanding of mom's mental health, which we know is critical to her child's well-being and development."

The researchers analyzed data from the Upstate KIDS study, which included babies born between 2008 and 2010 from 57 counties in New York State. The study followed 5,000 women for three years after their children were born.

Researchers assessed women's symptoms through a brief, five-item depression screening questionnaire, but the study did not clinically diagnose depression in the women. Women with underlying conditions, such as [mood disorders](#) and/or [gestational diabetes](#), were more likely to have higher levels of depressive symptoms that persisted throughout the study period.

The researchers noted that the study participants were primarily white,

non-Hispanic women. Future studies should include a more diverse, broad population to provide more inclusive data on [postpartum depression](#), Dr. Putnick said.

More information: Putnick, D, et al. Trajectories of Postpartum Depressive Symptoms. *Pediatrics*. September 2020.

Provided by NIH/Eunice Kennedy Shriver National Institute of Child Health and Human Development

Citation: Postpartum depression may persist three years after giving birth (2020, October 27) retrieved 11 May 2024 from <https://medicalxpress.com/news/2020-10-postpartum-depression-persist-years-birth.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--