


# Pulmonary artery thrombosis a complication of radiation therapy

October 9 2020

---



Credit: CC0 Public Domain

According to an article in ARRS' *American Journal of Roentgenology* (AJR), the imaging findings of in situ pulmonary artery thrombosis (PAT) associated with radiation therapy (RT) are different from those of

acute pulmonary emboli and do not appear to embolize. Due to the differences in clinical prognosis and subsequent management strategies, in situ PAT associated with RT—"which to our knowledge has not previously been described in the English literature," wrote the authors of this AJR article—must be distinguished from pulmonary embolism.

Searching the radiology database of a large teaching hospital to identify patients who had PAT develop after receiving RT, first author Jitesh Ahuja from the thoracic imaging department at the University of Texas' MD Anderson Cancer Center recorded the PAT's CT characteristics: number, location, appearance of filling defects, as well as the presence of associated lung fibrosis. The terminology (in situ thrombosis vs acute or chronic [pulmonary embolism](#)) used to describe PAT, the time between completion of RT and development of PAT, the size change of PAT, and any observation of new thrombi and emboli on follow-up imaging were also recorded.

With a study population consisting of 27 patients (19 men and eight women) at a mean age of 71 (range, 54-90 years), the primary malignancy was [lung cancer](#) in 22 patients (81%) and mesothelioma in five patients (19%). Whereas most PATs were solitary (93%) and nonocclusive (96%) and formed an obtuse angle to the vessel wall (89%), all PATs were eccentric within the involved pulmonary artery and located within the RT volume. The time from completion of RT to initial diagnosis of PAT on CT ranged from 53 to 2,522 days (mean, 675 days). In all patients, CT findings of radiation-induced lung fibrosis were present in the lung supplied by the affected pulmonary artery. "On follow-up imaging, none of the patients were observed to have filling defects develop in other parts of the PA, which would have suggested embolization," Ahuja et al. added.

Acknowledging that RT is a 'key choice' in multimodality treatment of intrathoracic malignancies, the authors noted that RT-associated

cardiovascular complications remain the leading non-cancer-related cause of morbidity and mortality among cancer survivors. "Radiologist awareness of PAT can facilitate accurate diagnosis and impact management," they concluded.

**More information:** Jitesh Ahuja et al, In Situ Pulmonary Artery Thrombosis: Unrecognized Complication of Radiation Therapy, *American Journal of Roentgenology* (2020). [DOI: 10.2214/AJR.19.22741](https://doi.org/10.2214/AJR.19.22741)

Provided by American Roentgen Ray Society

Citation: Pulmonary artery thrombosis a complication of radiation therapy (2020, October 9) retrieved 1 May 2024 from <https://medicalxpress.com/news/2020-10-pulmonary-artery-thrombosis-complication-therapy.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--