

Therapy plus medication better than medication alone in bipolar disorder

October 14 2020



Credit: CC0 Public Domain

A review of 39 randomized clinical trials by scientists from UCLA and their colleagues from other institutions has found that combining the use medication with psychoeducational therapy is more effective at

preventing a recurrence of illness in people with bipolar disorder than medication alone.

For the paper, published in *JAMA Psychiatry* researchers analyzed studies that included adult and adolescent patients currently receiving medication for [bipolar disorder](#) who were randomly assigned to either an active family, individual or [group therapy](#), or "usual care," meaning medication with routine monitoring and support from a psychiatrist.

David Miklowitz, Ph.D., the study's lead author, and a distinguished professor of psychiatry at the Jane and Terry Semel Institute for Neuroscience and Human Behavior at UCLA, said the studies reviewed followed patients for at least a year, measured rates of recurrence of bipolar disorder, depression and mania symptoms, and included study attrition or dropout rates.

The findings were:

- Psychoeducation with guided practice of illness management skills (for example, how to keep regular sleep and wake cycles) in a family or group format was more effective in reducing recurrences of mania and depressive symptoms than the same strategies in an individual therapy format.
- Cognitive [behavioral therapy](#), [family therapy](#) and interpersonal therapy were better at stabilizing [depressive symptoms](#) than other forms of treatment.
- Rates of dropout were lower in patients who received family-oriented therapies.
- Of the findings, Miklowitz said they point to the importance of having a support system.

"Not everyone may agree with me, but I think the family environment is very important in terms of whether somebody stays well," he said.

"There's nothing like having a person who knows how to recognize when you're getting ill and can say, 'you're starting to look depressed or you're starting to get ramped up.' That person can remind their loved one to take their medications or stay on a regular sleep-wake cycle or contact the psychiatrist for a medication evaluation."

Miklowitz said the same is true for a patient who may not have close relatives but does have support through group therapy.

"If you're in group therapy, other members of that group may be able to help you recognize that you're experiencing symptoms," he said. "People tend to pair off. It's a little bit like the AA model of having a sponsor."

More information: David J. Miklowitz et al, Adjunctive Psychotherapy for Bipolar Disorder, *JAMA Psychiatry* (2020). [DOI: 10.1001/jamapsychiatry.2020.2993](https://doi.org/10.1001/jamapsychiatry.2020.2993)

Provided by University of California, Los Angeles

Citation: Therapy plus medication better than medication alone in bipolar disorder (2020, October 14) retrieved 5 May 2024 from <https://medicalxpress.com/news/2020-10-therapy-medication-bipolar-disorder.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--