


# Nearly one in three young adults in the US does not know common stroke symptoms

October 26 2020

---



Credit: CC0 Public Domain

At a time when stroke is on the rise among young adults, nearly 30% of U.S. adults younger than age 45 do not know all five of the most common stroke symptoms, according to new research published today in *Stroke*, a journal of the American Stroke Association, a division of the American Heart Association.

Stroke is the No. 5 cause of death and a leading cause of disability in the United States. Each year, 10% to 15% of the nearly 795,000 people in the U.S. who have a stroke are young adults—between ages 18 and 45. Recent studies suggest [stroke incidence](#) is declining in the [general population](#), yet, stroke incidence and hospitalizations have increased by more than 40% in young adults in the past several decades.

"While the [medical community](#) has made significant improvements to reduce the severity and complications of strokes with early interventions, these efforts are of limited value if patients do not recognize stroke symptoms," said study author Khurram Nasir, M.D., M.P.H., M.Sc., chief of the division of cardiovascular prevention and wellness at Houston Methodist DeBakey Heart and Vascular Center in Houston, Texas. "Time is critical for treating stroke. The earlier people recognize symptoms, the better their chances are to reduce long-term disability from stroke."

To assess how well the U.S. population understands common stroke symptoms, Nasir and colleagues reviewed responses to the 2017 National Health Interview Survey. As part of the annual survey, adults are asked several questions about stroke including identifying five of the most common stroke symptoms, which in this survey were noted as:

- Numbness of face/arm/leg;
- Confusion/trouble speaking;
- Difficulty walking/dizziness/loss of balance;
- Trouble seeing in one/both eyes; and
- Severe headache.

In a targeted subset of the full survey, researchers analyzed answers from 9,844 younger adults, under age 45, which statistically represents 107.2 million young adults in the U.S. population. Average age of the younger respondents was 31, half were women and 62.2% were non-Hispanic

white.

The researchers found:

- Almost one in three (28.9%) respondents were not aware of all five common stroke symptoms.
- About 3% of respondents, representing nearly 3 million young adults, were not aware of any stroke [symptom](#).
- Hispanic adults and adults not born in the U.S. were about twice as likely to be unaware of any of the common stroke symptoms, compared to non-Hispanic White people and those born in the U.S.
- Young adults with a [high school diploma](#) or lower education level were nearly three-times as likely to be unaware of any stroke symptom, compared to young adults with higher education levels,

Nasir said the high number of young adults who remain unaware of stroke symptoms is surprising, and along with continued social inequities, these are major concerns.

"With the growing risk of stroke among younger adults in the U.S., our study sheds light on particularly vulnerable individuals and communities that already experience a disproportionately greater burden of stroke and cardiovascular risk factors, as well as reduced access to [health care services](#)," he said. "We hope that highlighting the continued impact of current health disparities may advance focused public health strategies and educational initiatives to increase awareness of and appropriate response to stroke symptoms."

The researchers also found that nearly 3% of young adults surveyed would not contact emergency medical services if they did see someone experiencing perceived stroke symptoms. "That finding could be a matter of life and death," said Mitchell S. V. Elkind, M.D., M.S.,

FAHA, FAAN, president of the American Heart Association.

"With proper, timely medical attention, stroke is largely treatable. The faster you are treated, the more likely you are to minimize the long-term effects of a stroke and even prevent death," said Elkind, professor of neurology and epidemiology at Vagelos College of Physicians and Surgeons and attending neurologist at New York-Presbyterian/Columbia University Irving Medical Center in New York City. "Calling 9-1-1 is critical because trained EMS personnel can start the care protocol en route to the hospital and have specialized teams standing by, ready at the hospital to administer the most appropriate treatment immediately."

Elkind says the American Heart Association/American Stroke Association advocates the use of the letters in ["F.A.S.T."](#) to spot stroke signs and to know when to call 9-1-1:

- Face drooping
- Arm weakness
- Speech slurred
- Time to call 9-1-1

Nasir said F.A.S.T. is among a number of creative and community-engaged initiatives that have aimed to increase public recognition of common [stroke symptoms](#), and he stresses an urgency to address contemporary health inequities head-on through such tailored and multidisciplinary public health approaches.

"Our results show that novel strategies are required at the population level to increase symptom awareness among [young adults](#), where we have found a higher-risk population with substantial variations in symptom recognition," he said.

A possible limitation of the study is that those surveyed responded with a

"yes" or "no" when asked if something was a common [stroke](#) symptom. This could lead to an overestimation of actual awareness rates, according to the authors.

**More information:** Reed Mszar et al. Association Between Sociodemographic Determinants and Disparities in Stroke Symptom Awareness Among US Young Adults, *Stroke* (2020). [DOI: 10.1161/STROKEAHA.120.031137](#)

Provided by American Heart Association

Citation: Nearly one in three young adults in the US does not know common stroke symptoms (2020, October 26) retrieved 30 April 2024 from <https://medicalxpress.com/news/2020-10-young-adults-common-symptoms.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--