

# Coming out as bisexual associated with increased risk of smoking: study

November 7 2020

---



Credit: CC0 Public Domain

For many years, lesbian, gay, bisexual, and other non-heterosexual (LGB+) folks have been known to be more likely to smoke than their straight counterparts.

But a new, first-of-its-kind Boston University School of Public Health (BUSPH) study paints a more precise picture by looking at LGB+ identities separately and over time, finding that bisexuality is the [identity](#) most associated with smoking, especially around the time of coming out.

Published in the journal *JAMA Pediatrics*, the nationally-representative cohort study followed 7,843 youth and [young adults](#) over three years, finding that those who came out as bisexual were twice as likely as consistently-heterosexual participants to start smoking. Coming out as lesbian, gay, or another non-heterosexual identity, or having a consistent LG+ identity, was not associated with being more likely to smoke.

The study "highlights the importance of moving beyond static measures of sexual identity towards more dynamic measures that capture critical periods of vulnerability," says Dr. Andrew Stokes, assistant professor of global health at BUSPH and the study's corresponding author.

"This approach turned out to be really important, because it revealed disparities that would have otherwise been missed if we measured identity at one time point, or grouped all LGB+ identities together," says study lead author Alyssa Harlow, a doctoral candidate at BUSPH.

"Bisexual young people may face unique forms of discrimination and stigma that increase their risk for smoking or other substance use behaviors," she says. "For example, they may experience stigma from heterosexual individuals as well as from within the LGB+ community. There's also prior research that shows that bisexual populations have worse mental health outcomes than LG+ populations.

"The findings point to a need for public health interventions specifically designed to address the unique needs, experiences, and stressors associated with coming out and identifying as bisexual," Harlow says.

For the study, the researchers used data from the first four waves of the nationwide Population Assessment of Tobacco and Health (PATH) study, which surveyed the same 14-29-year-olds three times between 2013 and 2018. (There were too few transgender respondents in this sample for the researchers to include gender identity in their analysis.) The researchers adjusted for other variables including sex, age, race/ethnicity, [education level](#) (for participants over 18) and parents' education level (for participants under 18), and where participants lived (urban/nonurban, and region of the U.S.).

By the third wave, 14% of the respondents had smoked at some point, and 6% were current smokers. The researchers found that the same sexual identity patterns held true both for having smoked at any point in the study period and for being a current smoker.

The researchers found that, compared to a consistent heterosexual identity, coming out as bisexual was associated with being more than twice as likely to smoke. Participants with LG+ identities in the first wave who shifted to a bisexual identity, or vice versa, were twice as likely to smoke.

On the other hand, participants with a consistent LG+ identity throughout the three waves of the study and participants who started out identifying as heterosexual and came out as LG+ were not more likely to smoke than those with a consistent heterosexual identity—while those with a consistent bisexual identity were slightly more likely to smoke.

The researchers say that the study's unique approach to LGB+ identities—separated and over time—could provide valuable insights for other issues that disproportionately affect the community, including mental health issues and substance use.

But to make that possible, more national surveys need to ask youth about

their sexual orientation and gender identity, says study co-author Dielle Lundberg, a research fellow at BUSPH.

"The PATH study is unique because it asks youth about their sexual orientation and gender identity. Most national surveys do not," Lundberg says.

"We must advocate for better data. Whenever national surveys fail to ask about sexual orientation and [gender identity](#), they are directly contributing to health inequities for LGBTQ+ populations."

**More information:** Alyssa F. Harlow et al. Association of Coming Out as Lesbian, Gay, and Bisexual+ and Risk of Cigarette Smoking in a Nationally Representative Sample of Youth and Young Adults, *JAMA Pediatrics* (2020). [DOI: 10.1001/jamapediatrics.2020.3565](https://doi.org/10.1001/jamapediatrics.2020.3565)

Provided by Boston University School of Medicine

Citation: Coming out as bisexual associated with increased risk of smoking: study (2020, November 7) retrieved 26 April 2024 from <https://medicalxpress.com/news/2020-11-bisexual.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--