

COVID-19 is making tinnitus worse, new study finds

November 6 2020



Credit: Pixabay/CC0 Public Domain

New research reveals that tinnitus, a common condition that causes the perception of noise in the ear and head, is being exacerbated by COVID-19—as well as the measures helping to keep us safe.

The study of 3,103 people with tinnitus was led by Anglia Ruskin University (ARU), with support from the British Tinnitus Association and the American Tinnitus Association. The study involved participants from 48 countries, with the vast majority coming from the UK and the US.

Published in the journal *Frontiers in Public Health*, the research found that 40% of those displaying symptoms of COVID-19 simultaneously experience a worsening of their tinnitus.

Although the study focused on people with pre-existing tinnitus, a small number of participants also reported that their condition was initially triggered by developing COVID-19 symptoms, suggesting that tinnitus could be a 'long COVID' symptom in some cases.

Tinnitus affects an estimated one in eight adults in the UK and is associated with reduced emotional wellbeing, depression, and anxiety.

The new study also found that a large proportion of people believe their tinnitus is being made worse by social distancing measures introduced to help control the spread of the virus. These measures have led to significant changes to work and lifestyle routines.

UK respondents reported this to be a greater issue compared to people from other countries, with 46% of UK respondents saying that [lifestyle changes](#) had negatively impacted their tinnitus compared to 29% in North America.

Internal worries such as fear of catching COVID-19, financial concerns, loneliness and trouble sleeping have contributed to making tinnitus more bothersome for 32% of people overall, with external factors such as increased videocalls, noisier home environments, home schooling and increased coffee and alcohol consumption also cited by respondents.

Females and the under-50s found tinnitus significantly more bothersome during the pandemic.

The study noted that as well as increasing the severity of tinnitus symptoms, the COVID-19 pandemic has also made it more difficult for people to access healthcare support for the condition. This could further increase emotional distress and worsen tinnitus symptoms, creating a vicious cycle. Before COVID-19, more than eight out of 10 UK patients were already unhappy with the treatment options available from their health professional.

Lead author Dr. Eldre Beukes, a Research Fellow at Anglia Ruskin University (ARU) in Cambridge, England, and Lamar University in Texas, said: "The findings of this study highlight the complexities associated with experiencing tinnitus and how both internal factors, such as increased anxiety and feelings of loneliness, and external factors, such as changes to daily routines, can have a significant effect on the condition.

"Some of the changes brought about by COVID-19 appear to have had a [negative impact](#) on the lives of people with tinnitus and participants in this study reported that COVID-19 symptoms are worsening or, in some cases, even initiating tinnitus and hearing loss. This is something that needs to be closely examined by both clinical and support services."

David Stockdale, Chief Executive of the British Tinnitus Association and a co-author of the study, said: "With the second wave of COVID-19 and the resulting national lockdown likely to increase feelings of stress and isolation, it's vital that we don't see the same mistakes as before when it comes to community health provision for people with tinnitus.

"Poor treatment of tinnitus in the early stages often leads to much worse cases and severe tinnitus can have a huge impact on mental health. With

this in mind, as the COVID-19 second wave takes hold, the healthcare system needs to ensure that anyone who develops [tinnitus](#) or experiences a worsening of their condition can access the professional healthcare support they need as quickly as possible."

More information: Eldré W. Beukes et al, Changes in Tinnitus Experiences During the COVID-19 Pandemic, *Frontiers in Public Health* (2020). [DOI: 10.3389/fpubh.2020.592878](https://doi.org/10.3389/fpubh.2020.592878)

Provided by Anglia Ruskin University

Citation: COVID-19 is making tinnitus worse, new study finds (2020, November 6) retrieved 25 April 2024 from <https://medicalxpress.com/news/2020-11-covid-tinnitus-worse.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--