

Study shows delirium can signal presence of COVID-19 in asymptomatic older patients in ED

November 19 2020



Credit: Pixabay/CC0 Public Domain

A study published today in *JAMA Network Open/Emergency Medicine* supports evidence that older persons admitted to emergency departments

(ED), and subsequently diagnosed positive for COVID-19, often present with delirium when they show no other typical COVID-19 symptoms, such as fever and cough. Sharon K. Inouye, M.D., M.P.H., Director of the Aging Brain Center in the Hinda and Arthur Marcus Institute for Aging Research at Hebrew SeniorLife and Professor of Medicine at Harvard Medical School, is senior author on the study.

Although COVID-19 poses a risk at all ages, adults aged 65 years and older are at greatest risk of severe disease, hospitalization, intensive care use, and death. Persons older than 65 years comprise 16 percent of the United States population yet have accounted for more than 80 percent of deaths in the U.S.

Delirium is an acute state of confusion, characterized by an altered level of consciousness, disorientation, inattention, and other cognitive disturbances. Beyond COVID-19, delirium is known to be a common symptom in [older adults](#) with severe disease in the ED and is associated with extended hospitalization, and increased morbidity and mortality. Despite the threat delirium poses to older ED patients, it is undetected in two-thirds of cases.

Researchers involved in the study examined 817 patients 65 or older admitted to the ED and who were diagnosed with COVID-19. They found almost a third had delirium at the time they were seen in the ED. A delirium diagnosis was the main presenting symptom for 16 percent of those patients, and 37 percent had no typical COVID-19 symptoms; delirium was the sixth most common presenting symptoms in all patients. These findings suggest the clinical importance of including delirium on checklists of presenting signs and symptoms of COVID-19 that guide screening, testing, and evaluation.

According to Dr. Inouye, "This study demonstrates that delirium is not only a common symptom of COVID-19, but also may be the leading and

possibly sole symptom in [older persons](#). Thus, [delirium](#) should be considered an important presenting symptom of COVID-19."

More information: Maura Kennedy, MD, MPH et al, Delirium in Older Patients With COVID-19 Presenting to the Emergency Department, *JAMA Netw Open*. 2020;3(11):e2029540. [DOI: 10.1001/jamanetworkopen.2020.29540](#)

Provided by Hebrew SeniorLife Hinda and Arthur Marcus Institute for Aging Research

Citation: Study shows delirium can signal presence of COVID-19 in asymptomatic older patients in ED (2020, November 19) retrieved 8 May 2024 from <https://medicalxpress.com/news/2020-11-delirium-presence-covid-asymptomatic-older.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--