

Extraction of largely-unexplored bodily fluid could be a new source of biomarkers

November 25 2020



Interstitial fluid (right) may provide an alternative source of biomarkers compared to blood (left) that could be useful in diagnosing human illness. Credit: Allison Carter, Georgia Tech


Using an array of tiny needles that are almost too small to see, researchers have developed a minimally-invasive technique for sampling a largely-unexplored human bodily fluid that could potentially provide a

new source of information for routine clinical monitoring and diagnostic testing.

Biochemical information about the body most commonly comes from analysis of blood—which represents only 6% of bodily fluids—but valuable information may also be found in other [bodily fluids](#) that are traditionally hard to get. Researchers have now developed a way to extract dermal interstitial [fluid](#) (ISF) - which circulates between cells in bodily tissues—using a simple through-the-[skin](#) technique that could provide a new approach for studying the metabolic products of cells, obtaining diagnostic biomarkers, and identifying potential toxins absorbed through the skin.

Because the dermal interstitial fluid doesn't clot like blood, the microneedle-based extraction could offer a new approach for continuous monitoring of glucose and other key health indicators.

Results of a human trial on the microneedle-based ISF sampling is reported Nov. 25 in the journal *Science Translational Medicine*. The study, conducted by researchers from the Georgia Institute of Technology and Emory University, was supported in part by the National Institutes of Health.




Interstitial fluid (right) may provide an alternative source of biomarkers compared to blood (left) that could be useful in diagnosing human illness. Credit: Allison Carter, Georgia Tech

"Interstitial fluid originates in the blood and then leaks out of capillaries to bring nutrients to cells in the body's tissues. Because interstitial fluid is in direct communication with the cells, it should have information about the tissues themselves beyond what can be measured from testing the blood," said Mark Prausnitz, Regents' Professor and J. Erskine Love Jr. Chair in Georgia Tech's School of Chemical and Biomolecular Engineering, "This microneedle-based technique could provide a minimally-invasive and simple way to access this interstitial fluid to make it available for medical diagnostic and research applications."

ISF has been difficult to sample. Indwelling instruments for monitoring glucose in ISF already exist, and other researchers have used surgically-implanted tubing and vacuum-created blisters to extract ISF through the skin, but these techniques are not suitable for routine clinical diagnostic use.

The researchers, led by first author Pradnya Samant, used a patch containing five solid stainless-steel microneedles that were a hundredth of an inch in length. By pressing the patch at an angle into the skin of 50 human subjects, they created shallow micropores that reached only into the outer layer of skin containing ISF. The researchers then applied a suction to the area of skin containing the pores and obtained enough ISF to do three types of analysis. For comparison, they also took [blood samples](#) and obtained ISF using the older blister technique.

To accurately determine the biomarkers available in the ISF, the researcher needed to avoid getting blood mixed with the ISF. Though major blood vessels don't exist in the outer layers of skin, capillaries there can be damaged by the insertion of the microneedles. In their studies, the researchers found that if they slowly ramped up the suction after inserting the microneedles, they could obtain fluid clear of blood.



Microneedle patches containing micron-scale needles are used to create temporary pores in the skin through which interstitial fluid can be extracted. Credit: Allison Carter, Georgia Tech

The overall extraction procedure took a total of about 20 minutes for each test subject. The procedure was well tolerated by the volunteers, and the microscopic pores healed quickly within a day with minimal irritation.

The extracted fluid was analyzed at Emory University using liquid chromatography-mass spectrometry techniques to identify the chemical species it contained. Overall, there were about 10,000 unique compounds, most of which were also found in the blood samples.

However, about 12 percent of the chemical species were not found in the blood, and others were found in the ISF at higher levels than in the blood.

"The skin is metabolically active, and it is full of cells that are changing the fluid," Prausnitz said. "We found that some of the compounds were unique to the ISF, or enriched there, and that is what we were hoping to find."

While not all the compounds unique to the ISF could be analyzed, the research team identified components of products that are applied to the skin—such as hand lotions—and pesticides that may enter the body through the skin. This discovery could set the stage for use of the microneedle technique for dermatological and toxicology studies.

"If you want to look at what accumulates in the skin over time, this may provide a way to obtain information about those kinds of exposures," Prausnitz said. "These are materials that may accumulate in the tissues of our body, but are not found in the bloodstream."

The researchers also determined the pharmacokinetics of caffeine and the pharmacodynamics of glucose—both [small molecules](#)—from the ISF, indicating that that dynamic biomarker information could be obtained from the technique. Those measurements suggested that ISF could provide a means for continuously monitoring of such compounds, taking advantage of the fact that the fluid does not clot.

"We were encouraged that we found a good correlation between the [blood](#) and interstitial fluid glucose, which suggests we might be able to have a continuous glucose monitoring system based on this technology," Prausnitz said. A microneedle-based system could provide a less-invasive alternative to existing implantable glucose sensors by allowing the sensing components to remain on the surface of the skin.

In future research, Prausnitz would like to reduce the time required to extract the ISF and simplify the process by eliminating the vacuum pump. Additional study of the compounds found in the fluid could also show whether they may have medical diagnostic value.

"We'd like to make this microneedle-based technique available to the research community to make ISF routinely available for study," he said. "Tissue interstitial fluid could be a novel source of biomarkers that complements conventional sources. This research provides a means to study this further."

More information: P.P. Samant et al., "Sampling interstitial fluid from human skin using a microneedle patch," *Science Translational Medicine* (2020). [stm.sciencemag.org/lookup/doi/ ... scitranslmed.aaw0285](https://stm.sciencemag.org/lookup/doi/10.1126/scitranslmed.aaw0285)

Provided by Georgia Institute of Technology

Citation: Extraction of largely-unexplored bodily fluid could be a new source of biomarkers (2020, November 25) retrieved 25 April 2024 from <https://medicalxpress.com/news/2020-11-largely-unexplored-bodily-fluid-source-biomarkers.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--