


Loneliness in Parkinson's disease may lead to worsening of symptoms

November 18 2020



Immunohistochemistry for alpha-synuclein showing positive staining (brown) of an intraneural Lewy-body in the Substantia nigra in Parkinson's disease. Credit: Wikipedia

Research from UCLA scientists and colleagues from other institutions

finds that people with Parkinson's disease who lack meaningful social interactions may be at an increased risk for severe symptoms related to the disease.

The study, which was published in the journal *NPJ Parkinson's Disease*, evaluated not only the social and emotional life of patients, but also their nutrition and exercise habits.

Over a 5-year period, from 2014-2019, researchers at UCLA, the University of Washington and Bastyr University collected information from 1,500 people with Parkinson's disease. Study participants were surveyed most recently in December. Participants who reported being most lonely, also reported exercising less, were less likely to follow healthier diets and experienced a lower quality of life.

"That surprised us," says study author Dr. Indu Subramanian, a neurologist at the David Geffen School of Medicine at UCLA and director of the Southwest Veteran Affairs Parkinson's Disease Research, Education and Clinical Centers.

"One of the most detrimental things is actually being lonely," Subramanian says. The negative impact of loneliness on symptom severity, she says, was as large as the positive effect from exercise.

Millions of people have curtailed their social interactions to stay safe during the COVID-19 pandemic. While this is good practice for avoiding a virus, the resulting isolation and loneliness can create a health hazard that could be particularly hard on people with Parkinson's disease who sometimes limit interactions due to adverse disease symptoms such as tremor.

Subramanian and the rest of the team recently sent out a new survey to the study participants to collect data about how the pandemic has

affected their symptoms.

Even patients who have happy family lives can suffer from loneliness, she adds, though that may seem counterintuitive.

People thrive in three spheres of [social interaction](#): One is the intimate connection of a marriage or partnership; the next wider sphere is a circle of friends; and the third is belonging to a group with a shared sense of identity.

"For people with Parkinson's disease, they may be in a support group. It could even be something like a book club," Subramanian says.

To support that shared sense of identity during the pandemic, Subramanian started a virtual support group, which meets two to three times a week, for people with Parkinson's disease.

"It's called social prescribing, because you literally prescribe to your patients to be more socially connected," Subramanian says. "It's actually grown into an international group of patients. People have grown to enjoy and connect through it. Some people have told me it's the only social thing they do at all."

More information: Indu Subramanian et al, Synergy of pandemics-social isolation is associated with worsened Parkinson severity and quality of life, *npj Parkinson's Disease* (2020). [DOI: 10.1038/s41531-020-00128-9](https://doi.org/10.1038/s41531-020-00128-9)

Provided by University of California, Los Angeles

Citation: Loneliness in Parkinson's disease may lead to worsening of symptoms (2020, November

18) retrieved 18 April 2024 from <https://medicalxpress.com/news/2020-11-loneliness-parkinson-disease-worsening-symptoms.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.