


Lovestruck by oxytocin. Novel roles of the hormone in controlling male sexual function

November 18 2020



Oxytocin directly activates the spinal ejaculation generator (SEG)/gastrin-releasing peptide (GRP) neurons via oxytocin receptors (OXTRs) and influences male sexual function in the rat lumbar spinal cord. Release of oxytocin in the lumbar cord is not limited to conventional synapses and acts by diffusion—a localized volume transmission—to reach OXTRs on SEG/GRP neurons and facilitate male sexual activity. Credit: 2020 Okayama University

Hormones are key players of the endocrine system and have a major influence on our emotional and sexual wellbeing. The hormone oxytocin is involved in a wide range of emotions from social bonding to maternal behaviors like nursing and lactation. But the most popular and well-known role of oxytocin, lending it its popular moniker of 'love hormone,' is its role in romantic and sexual emotions.

The functional mechanism of [oxytocin](#) in male sexual function and behavior is not clearly understood, but there is some evidence supporting the role of oxytocin-specific nerve cells or neurons in the brain that project to the lower spinal cord and control penile erection and ejaculation in male rats. Now, in a brand new study published in *Current Biology*, a group of researchers led by Professor Hirotaka Sakamoto from Okayama University, Japan, has explored this potential role of oxytocin and the underlying mechanisms in modulating male sexual function using rats as a model system.

Oxytocin is transferred from the brain to various parts of the body by the blood, and from neuron to neuron through structures called "synapses." However, the precise mechanisms by which sparsely dispersed oxytocin fibers—structures responsible for responding to oxytocin in the central nervous system—cause the activation of widely distributed receptors remain unclear.

The researchers from Japan investigated a novel non-synaptic mode of oxytocin transport across the central nervous system. When asked to explain this process, Prof Sakamoto refers to an interesting analogy: "Overall, the endocrine system, which acts on widespread distant organs via the circulation, resembles a 'broadcasting satellite' communication, whereas [synaptic transmission](#) resembles the hard-wired 'ethernet.' Accordingly, the localized volume transmission of peptides resembles

'Wi-Fi' communication, since it is a hybrid of both endocrine (satellite) and synaptic (ethernet) systems, and may be the predominant mechanism of oxytocinergic modulation of socio-sexual behavior and cognition throughout the central nervous system."

It is already known that spinal regions like the spinal ejaculation generator (SEG) are known to control sexual functions in male rodents. To assess the role of oxytocin in copulatory and ejaculatory responses, the team injected oxytocin into the spine of male rats. The gastrin-releasing peptide or GRP neurons are an important component of the SEG, as they control the lower lumbar region connected to muscles at the base of the penis, thereby controlling erection and ejaculation.

Oxytocin caused an increased sexual activity and neuronal activity in injected animals. More specifically, oxytocin was found to directly activate SEG/GRP neurons via oxytocin receptors, which detect oxytocin, and influence male sexual function in the rat lumbar spinal cord. Using an oxytocin receptor antagonist, which reduces the activity of oxytocin receptors, resulted in a latency and decrease in number of [sexual activity](#) and ejaculatory responses in majority of the animals, confirming the importance of oxytocin.

The question remained about the transport of oxytocin, however. Electron microscopy images acquired from slices of the lumbar region ruled out the presence of synaptic vesicles or connections. Upon stimulation of exocytosis ex-vivo, they were able to observe oxytocin transport mediated by a more passive diffusion in extracellular spaces at non-synaptic sites.

Highlighting the importance of the study, Prof Sakamoto remarks, "Now that we have uncovered a novel neural mechanism-the 'localized volume transmission' of oxytocin from axons-involved in controlling male sexual function in the spinal cord, we can hope that this may lead to the

development of treatments for male sexual dysfunction."

This study thus presents a completely unprecedented role for oxytocin in male sexual function, in addition to its long-standing "female-centric" role. Learning more about this "love hormone" may indeed help us foster healthier and long-lasting loving relationships!

More information: Takumi Oti et al, Oxytocin Influences Male Sexual Activity via Non-synaptic Axonal Release in the Spinal Cord, *Current Biology* (2020). [DOI: 10.1016/j.cub.2020.09.089](https://doi.org/10.1016/j.cub.2020.09.089)

Provided by Okayama University

Citation: Lovestruck by oxytocin. Novel roles of the hormone in controlling male sexual function (2020, November 18) retrieved 3 May 2024 from <https://medicalxpress.com/news/2020-11-lovestruck-oxytocin-roles-hormone-male.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--