


Patients at risk of atrial fibrillation may need additional monitoring after heart surgery

November 16 2020



A 12 lead ECG showing atrial fibrillation at approximately 150 beats per minute.
Credit: James Heilman, MD/Wikipedia/CC BY-SA 3.0

Atrial fibrillation (AF), an irregular heartbeat that can increase the likelihood of stroke, was detected up to 10 times more frequently in high risk patients recovering from heart surgery who wore a continuous cardiac monitor for a month, compared to patients who had usual follow-up care following their procedure, according to late breaking research presented today at the American Heart Association's Scientific Sessions 2020.

"The incidence of post-operative atrial fibrillation (POAF) after

discharge from [cardiac surgery](#) is not well defined," said Subodh Verma, M.D., Ph.D., a cardiac surgeon at St. Michael's Hospital, University of Toronto in Canada, and one of the lead authors of the study. "Most studies are limited to the hospitalization phase only; studies beyond hospitalization are few. In addition, very little is known about patients who have little to no AF during hospitalization after cardiac [surgery](#), and they are often sent home with no treatment. Therefore, the question of whether the risk of POAF extends after hospitalization remains an important and unanswered question, especially for patients who had no previous history of AF. Quantifying the risk of ongoing POAF after hospitalization is important because many patients have elevated stroke risk scores that increase the need for prompt AF treatment."

Researchers followed a total of 336 patients treated at eight medical centers across Canada who were recovering from cardiac surgery. The study participants did not have a history of atrial fibrillation prior to surgery and were at-risk for stroke based upon factors including their cardiac health, age and medical history.

The patients in the study were randomly assigned to two groups: 163 study participants underwent 30 days of 24-hour, real time, cardiac rhythm monitoring through an adhesive, patch-based monitor worn on their chest; 173 participants were in the [control group](#) and received 30 days of usual care, which did not involve planned cardiac rhythm testing/electrocardiogram assessment unless it was deemed medically necessary. The patients in both groups then had 14 days of continuous cardiac rhythm monitoring approximately six to nine months after their surgery.

The study determined that in the patient group who wore the 24-hour monitoring patch, [atrial fibrillation](#) was detected at a rate 10 times higher than in those who received usual care. Atrial fibrillation or [atrial flutter](#) lasting ≥ 6 minutes occurred and was detected in 32 (19.6%) patients

from the monitoring group and in three (1.7%) patients who received normal care during the 30 days they were followed.

"Our study points to the fact that POAF is not self-limited to hospital stay per se. A significant risk of POAF persists even in those patients without any preoperative or pre-discharge AF. These data may help inform physicians about the importance of surveillance and vigilance in patients at high risk of stroke with respect to monitoring and prompt treatment for AF," added Verma.

More information: [www.abstractsonline.com/pp8/? ...
#!/9144/session/1456](http://www.abstractsonline.com/pp8/?...#!/9144/session/1456)

Provided by American Heart Association

Citation: Patients at risk of atrial fibrillation may need additional monitoring after heart surgery (2020, November 16) retrieved 17 April 2024 from <https://medicalxpress.com/news/2020-11-patients-atrial-fibrillation-additional-heart.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--