

Rapid point-of-care testing during and after COVID-19 – how widely should it be used?

November 18 2020



Credit: Pixabay/CC0 Public Domain

Even before the COVID-19 pandemic, the point-of-care testing industry was investing millions of pounds to develop rapid tests to tell us the cause of respiratory infections. The pandemic has accelerated this process. In an editorial published in the *British Journal of General Practice* today [17 November], researchers from the University of Bristol's Centre for Academic Primary Care ask if we know enough

about these tests to merit their widespread use in primary care.

The editorial suggests that like a new medicine or vaccine, the safety, clinical and cost-effectiveness of rapid point-of-care tests should be investigated in carefully conducted [clinical trials](#) before they are introduced into widespread use.

Alastair Hay, Professor of Primary Care in Bristol Medical School: Population Health Sciences (PHS) and co-author, said: "Rapid point-of-care tests are attractive, because traditional laboratory methods take hours to days, often too long for clinical decision making. They can deliver results in less than one hour, so what's not to like?"

"The main issue is we do not understand if they are accurate enough, effective enough or cost-effective enough to merit widespread introduction. This is not a problem, because accuracy, effectiveness and [cost-effectiveness](#) can be investigated—just as we would a new medicine or vaccine.

"Rapid point-of-care tests have been advocated as key to future antimicrobial stewardship—a way of decreasing the diagnostic uncertainty which has been shown to be a leading cause of antibiotic overprescribing.

"There is great potential for these tests to be an important part of the armory in the fight against antimicrobial resistance, but we have a responsibility to ensure that they are safe and reliable before investing large sums of public money in their use."

More information: Hannah V Thornton et al. Point-of-care testing for respiratory infections during and after COVID-19, *British Journal of General Practice* (2020). [DOI: 10.3399/bjgp20X713561](https://doi.org/10.3399/bjgp20X713561)

Provided by University of Bristol

Citation: Rapid point-of-care testing during and after COVID-19 – how widely should it be used? (2020, November 18) retrieved 26 April 2024 from <https://medicalxpress.com/news/2020-11-rapid-point-of-care-covid-widely.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.