

# Team makes breakthrough discovery on brain cortex functionality

November 11 2020, by Lauren Moriarty

---



Credit: CC0 Public Domain

A team of researchers from UTSA's Neurosciences Institute is challenging the historical belief that the organization of the cortical circuit of GABAergic neurons is exclusively local.

UTSA College of Sciences researchers Alice Bertero, Hector Zurita and Marc Normandin and biology associate professor Alfonso Junior

Apicella collaborated on the research project.

In the past, labeling [individual neurons](#) allowed researchers to study the [neurons](#) that project from the brain's [cortex](#) to the striatum. The results suggested that the pathways are exclusively excitatory. For this reason most people assumed that inhibition must occur when excitatory cortical neurons activate intrastriatal inhibitory.

The UTSA team contested this view by providing anatomical and physiological evidence for the existence of long-range parvalbumin-expression neurons from the cortex to the striatum in the brains of mice.

This finding is an essential step toward better understanding the neuronal mechanism of cortical long-range GABAergic neurons in healthy and diseased brains. In particular this could lead to new treatments for patients with epilepsy, post-[traumatic stress disorder](#), schizophrenia and other mental conditions in which GABAergic neurons play a significant role.

"Daily we are bombarded by multiple [auditory stimuli](#), including various speeches," said Apicella. "Particularly, listeners must determine which speech acoustic features are relevant and generalize across irrelevant ones during speech perception. We think that the interaction between the cortex and the striatum is fundamental to optimize nonnative speech learning in adulthood. Therefore, it is crucial to determine which of the cortical-striatum pathway's cellular elements are involved in this process."

The discovery that the striatum receives both excitatory and inhibitory inputs from the cortex suggests that these inputs' timing and relative strength can modulate the striatum's activity. This is important because it invites the [scientific community](#) to speculate that the long-range parvalbumin-expressing projects could play a role through gamma

oscillation synchronization between the auditory cortex and striatum.

The team proposes that future experiments will provide further insight into the role of the timing and ratio of excitation and inhibition, two opposing forces in the mammalian [cerebral cortex](#), affecting the cortico-striatal dynamic.

"Future studies," said Apicella, "will provide a general mechanism of cortico-striatal oscillation involved in reward learning and action-selection behavior driven by the auditory stimuli."

**More information:** Alice Bertero et al, Auditory Long-Range Parvalbumin Cortico-Striatal Neurons, *Frontiers in Neural Circuits* (2020). [DOI: 10.3389/fncir.2020.00045](https://doi.org/10.3389/fncir.2020.00045)

Provided by University of Texas at San Antonio

Citation: Team makes breakthrough discovery on brain cortex functionality (2020, November 11) retrieved 26 April 2024 from <https://medicalxpress.com/news/2020-11-team-breakthrough-discovery-brain-cortex.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--