

Technology lets clinicians objectively detect tinnitus for first time

November 18 2020



A person set up for an fNIRS recording (the brain imaging technique used in this paper). Credit: Mehrnaz Shoushtarian



A technology called functional near-infrared spectroscopy (fNIRS) can be used to objectively measure tinnitus, or ringing in the ears, according to a new study published November 18 in the open-access journal *PLOS ONE* by Mehrnaz Shoushtarian of The Bionics Institute, Australia, and colleagues.

Tinnitus, the perception of a high-pitched ringing or buzzing in the ears, affects up to 20% of adults and, when severe, is associated with depression, cognitive dysfunction and stress. Despite its wide prevalence, there has been no clinically-used, objective way to determine the presence or severity of tinnitus.

In the new study, researchers turned to fNIRS, a non-invasive and nonradioactive imaging method which measures changes in blood oxygen levels within <u>brain tissue</u>. The team used fNIRS to track activity in areas of the brain's cortex previously linked to tinnitus. They collected fNIRS data in the resting state and in response to auditory and <u>visual stimuli</u> in 25 people with chronic tinnitus and 21 controls matched for age and hearing loss. Participants also rated the severity of their tinnitus using the Tinnitus Handicap Inventory.

fNIRS revealed a statistically significant difference in the connectivity between areas of the brain in people with and without tinnitus. Moreover, the brain's response to both visual and <u>auditory stimuli</u> was dampened among patients with tinnitus. When a <u>machine learning</u> approach was applied to the data, a program could differentiate patients with slight/mild tinnitus from those with moderate/severe tinnitus with an 87.32% accuracy. The authors conclude that fNIRS may be a feasible way to objectively assess tinnitus to assess new treatments or monitor the effectiveness of a patient's treatment program.

The authors add: "Much like the sensation itself, how severe an individual's tinnitus is has previously only been known to the person



experiencing the condition. We have combined machine learning and non-invasive brain imaging to quantify the severity of tinnitus. Our ability to track the complex changes that <u>tinnitus</u> triggers in a sufferer's brain is critical for the development of new treatments."

More information: Shoushtarian M, Alizadehsani R, Khosravi A, Acevedo N, McKay CM, Nahavandi S, et al. (2020) Objective measurement of tinnitus using functional near-infrared spectroscopy and machine learning. *PLoS ONE* 15(11): e0241695. doi.org/10.1371/journal.pone.0241695

Provided by Public Library of Science

Citation: Technology lets clinicians objectively detect tinnitus for first time (2020, November 18) retrieved 3 May 2024 from <u>https://medicalxpress.com/news/2020-11-technology-clinicians-tinnitus.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.