

## Vitamin C's effectiveness against COVID may hinge on vitamin's natural transporter levels

November 11 2020



Drs. Sadanand Fulzele and Carlos Isales. Credit: Kim Ratliff, Production Coordinator, Augusta University

High doses of vitamin C under study for treating COVID-19 may benefit some populations, but investigators exploring its potential in aging say



key factors in effectiveness include levels of the natural transporter needed to get the vitamin inside cells.

Age, race, gender, as well as <u>expression levels</u> and genetic variations of those vitamin C transporters that make them less efficient, all may be factors in the effectiveness of vitamin C therapy against COVID-19 and other maladies, investigators at the Medical College of Georgia Center for Healthy Aging report in a commentary in the journal *Aging and Disease*.

The investigators recommend that those factors be considered in the design and execution of clinical trials, and when trial results are analyzed, for COVID-19 as well as other conditions, says Dr. Sadanand Fulzele, aging researcher and the article's corresponding author.

The novel nature and lack of immunity against the coronavirus has prompted a worldwide pursuit of effective treatments for COVID-19, they write. That includes repurposing drugs with known safety profiles, including Vitamin C, an established <u>immune system booster</u> and antioxidant, which made it a logical choice to explore in COVID-19. Both strategies are needed in response to infection with the novel coronavirus to ensure a strong immune response to stop the virus from replicating in the body, and to avoid the over-the-top, destructive immune response the virus itself can generate if it does.

There are at least 30 clinical trials underway in which vitamin C, alone or in combination with other treatments, is being evaluated against COVID-19, some with doses up to 10 times the recommended 65 to 90 milligrams daily of vitamin C.

Factors like whether or not vitamin C can get inside the cell, likely are an issue in the effectiveness the therapies ultimately show, says Dr. Carlos M. Isales, co-director of the MCG Center for Healthy Aging and



chief of the MCG Division of Endocrinology, Diabetes and Metabolism.

In fact, without adequate transporters on a cell's surface to get the watersoluble vitamin past the lipid layer of cell membranes, particularly large doses may enable the vitamin to cluster around the outside of cells where it actually starts producing oxidants, like damaging reactive oxygen species, rather than helping eliminate them, says Isales, a study coauthor.

"We think it's important to look at transporter expression," Fulzele says.

They suspect low transporter expression is a factor in the mixed results from vitamin C's use in a variety of other conditions. Clinical trials in osteoarthritis, for example, an autoimmune disease where a misdirected immune system is attacking the joints, has gotten mixed results, Fulzele says. However its usage in other viral-induced problems, like potentially deadly sepsis, has shown benefit in reducing organ failure and improving lung function in <u>acute respiratory distress syndrome</u>, which is also a major cause of sickness and death with COVID-19.

At the time their *Aging and Disease* paper was published, there were not yet published studies of the efficacies of high-dose, intravenous vitamin studies underway for COVID-19.

Fulzele, who works on vitamin C in aging, and others have shown that some conditions, like osteoarthritis and even normal aging, are associated with significant downregulation of at least one subtype of vitamin C transporter.

In fact, part of the paradox and concern with COVID-19 is that those most at risk mostly have both lower levels of vitamin C before they get sick and fewer transporters to enable the vitamin to be of benefit if they get more, Fulzele says.



Many of those most at risk from COVID-19, including individuals who are older, Black, male and with chronic medical conditions like osteoarthritis, hypertension and diabetes, tend to have lower levels of vitamin C, another reason vitamin C therapy would be considered a reasonable treatment, Isales says. The investigators also note that patients may develop a vitamin C deficiency over the course of their COVID-19 illness since, during an active infection, vitamin C is consumed at a more rapid rate. Insufficient levels can augment the damage done by an overzealous immune response.

While not routinely done, transporter expression can be measured today using PCR technology, a method also used for novel <u>coronavirus</u> as well as influenza testing. While increasing transporter expression is not yet doable in humans, one of Fulzele's many research goals is to find a drug or other method to directly increase expression, which should improve the health of older individuals as well as those with other medical conditions that compromise those levels.

He notes that reduced transporter levels that occur naturally with age are a factor in the reduced immune function that also typically accompanies aging. That means that even when a 60-year-old and 20-year-old both have a healthy diet in which they consume similar, sufficient amounts of vitamin C, the vitamin is not as effective at boosting the older individual's immune response. Reduced immune function in older individuals is known to put them at increased risk for problems like cancer and COVID-19.

Low vitamin C levels also have been correlated with higher mortality in older individuals from causes like cardiovascular disease. High oxidative stress, a major factor in conditions like cardiovascular disease as well as aging and now COVID-19, also is associated with significantly reduced expression of the vitamin C transporter.



Isales and Fulzele doubt that taking a lot of vitamin C is a good preventive strategy against COVID-19, except in those individuals with a known deficiency.

Vitamin C is an essential vitamin, which means people have to consume it in their food or supplements. Foods naturally high in vitamin C include oranges, potatoes, tomatoes, broccoli and Brussels sprouts. The <u>vitamin</u>'s diverse roles in the body also include formation of blood vessels, collagen and cartilage.

**More information:** Gregory Patterson, Carlos M. Isales, Sadanand Fulzele. Low level of Vitamin C and dysregulation of Vitamin C transporter might be involved in the severity of COVID-19 Infection[J]. Aging and disease,

www.aginganddisease.org/EN/10.14336/AD.2020.0918

## Provided by Medical College of Georgia at Augusta University

Citation: Vitamin C's effectiveness against COVID may hinge on vitamin's natural transporter levels (2020, November 11) retrieved 6 May 2024 from <u>https://medicalxpress.com/news/2020-11-vitamin-effectiveness-covid-hinge-natural.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.