

Brain imaging predicts PTSD after brain injury

December 29 2020

Posttraumatic stress disorder (PTSD) is a complex psychiatric disorder brought on by physical and/or psychological trauma. How its symptoms, including anxiety, depression and cognitive disturbances arise remains incompletely understood and unpredictable. Treatments and outcomes could potentially be improved if doctors could better predict who would develop PTSD. Now, researchers using magnetic resonance imaging (MRI) have found potential brain biomarkers of PTSD in people with traumatic brain injury (TBI).

The study appears in *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, published by Elsevier.

"The relationship between TBI and PTSD has garnered increased attention in recent years as studies have shown considerable overlap in risk factors and symptoms," said lead author Murray Stein, MD, MPH, FRCPC, a Distinguished Professor of Psychiatry and Family Medicine & Public Health at the University of California San Diego, San Diego, La Jolla, CA, USA. "In this study, we were able to use data from TRACK-TBI, a large longitudinal study of patients who present in the Emergency Department with TBIs serious enough to warrant CT (computed tomography) scans."

The researchers followed over 400 such TBI patients, assessing them for PTSD at 3 and 6 months after their [brain injury](#). At 3 months, 77 participants, or 18 percent, had likely PTSD; at 6 months, 70 participants or 16 percent did. All subjects underwent brain imaging after injury.

"MRI studies conducted within two weeks of injury were used to measure volumes of key structures in the brain thought to be involved in PTSD," said Dr. Stein. "We found that the volume of several of these structures were predictive of PTSD 3-months post-injury."

Specifically, smaller volume in [brain regions](#) called the cingulate cortex, the superior frontal cortex, and the insula predicted PTSD at 3 months. The regions are associated with arousal, attention and emotional regulation. The structural imaging did not predict PTSD at 6 months.

The findings are in line with previous studies showing smaller volume in several of these brain regions in people with PTSD and studies suggesting that the reduced cortical volume may be a risk factor for developing PTSD. Together, the findings suggest that a "brain reserve," or higher cortical volumes, may provide some resilience against PTSD.

Although the biomarker of brain volume differences is not yet robust enough to provide clinical guidance, Dr. Stein said, "it does pave the way for future studies to look even more closely at how these brain regions may contribute to (or protect against) mental health problems such as PTSD."

Cameron Carter, MD, Editor of *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, said of the work, "This very important study uses [magnetic resonance](#) imaging to take the field a step closer to understanding why some people develop PTSD after trauma and others do not. It also lays the groundwork for future research aimed at using [brain](#) imaging to help predict that a person is at increased risk and may benefit from targeted interventions to reduce the clinical impact of a traumatic event."

More information: Murray B. Stein et al, Smaller Regional Brain Volumes Predict Posttraumatic Stress Disorder at 3 Months after Mild

Traumatic Brain Injury, *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging* (2020). [DOI: 10.1016/j.bpsc.2020.10.008](https://doi.org/10.1016/j.bpsc.2020.10.008)

Provided by Elsevier

Citation: Brain imaging predicts PTSD after brain injury (2020, December 29) retrieved 26 April 2024 from <https://medicalxpress.com/news/2020-12-brain-imaging-ptsd-injury.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.