


Chatbot may detect early dementia in time for intervention

December 7 2020



Credit: Unsplash/CC0 Public Domain

If Alzheimer's dementia is identified early, at the stage of Mild Cognitive Impairment, for instance, the decline in neural functioning can be stabilized or even curtailed in some cases, research has found.

QUT Ph.D. data science researcher Ahmed Alkenani has developed automatic machine learning models using language features to identify multiple stages of dementia, including Mild Cognitive Impairment (MCI), possible Alzheimer's dementia (PoAD), and probable Alzheimer's dementia (AD), ie not fully developed AD.

"It is possible to identify language changes years prior to developing dementia, which highlights the importance of linguistic analysis for early dementia detection," said Mr Alkenani, from QUT's School of Computer Science.

"Our study shows that early stages of dementia can be efficiently diagnosed through linguistic patterns and deficits using machine learning models.

"Early, accurate diagnosis is important to enable clinicians to intervene in time to delay or prevent Alzheimer's dementia.

"Currently the initial diagnosis is performed with pen-and-paper screening tests such as the Mini Mental State Examination (MMSE) and the Montreal Cognitive Assessment (MoCA).

"These traditional tests are normally clinic-based and involve a series of questions and tasks to assess short-term memory, attention, repetition and orientation.

"Traditional tests rely on the neurologist's experience and level of expertise for delivery and assessment and the results are typically affected by the patient's age (possibility of normal age-related cognitive decline) and level of education."

Mr Alkenani said that as dementia advanced a person's language comprehension and spoken complexity declined.

"Dementia severity is associated with a limited vocabulary and increased word repetitions giving patterns we can pick up as linguistic biomarkers as dementia progresses," Mr Alkenani said.

"Our study analyzed language samples from DementiaBank, a large open-source database of language samples from people with various stages of cognitive impairment and dementia who took the well-known 'Cookie Theft' picture description task.

"For this study we introduced several new word and grammar features alongside previously established ones to train machine learning classifiers to identify linguistic biomarkers of MCI and AD.

"We studied 236 language samples from people diagnosed with probable AD, 43 samples from people with MCI, 21 samples belonging to people with possible AD, and 243 from healthy people.

"We found people with dementia leaned towards using fewer nouns but more verbs, pronouns, and adjectives as dementia progressed compared to healthy adults.

"For instance, we found noun to verb ratio and verb to noun ratio to be significant in differentiating both AD and MCI from healthy people.

"This is interesting, as previous research has revealed that nouns and verbs are learnt and activated in different brain regions which could be matched with the area of the brain that is first affected by dementia and help [early intervention](#)."

Mr Alkenani said the study was believed to be the first to classify AD, MCI and PoAD accurately and automatically through machine learning models.

"We explored word and syntax patterns and we correlated these patterns across all stages to highlight the strongest associations.

"Our ultimate aim is to develop a conversational agent or chatbot that could be used remotely to facilitate the initial diagnosis of early stage [dementia](#) as an attempt to replace traditional screening tests."

"Predicting Prodromal Dementia Using Linguistic Patterns and Deficits" was published in *IEEE Access*.

More information: Ahmed H. Alkenani et al. Predicting Prodromal Dementia Using Linguistic Patterns and Deficits, *IEEE Access* (2020). [DOI: 10.1109/ACCESS.2020.3029907](https://doi.org/10.1109/ACCESS.2020.3029907)

Provided by Queensland University of Technology

Citation: Chatbot may detect early dementia in time for intervention (2020, December 7) retrieved 9 April 2024 from <https://medicalxpress.com/news/2020-12-chatbot-early-dementia-intervention.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--