


# Stroke in childhood and adolescence: Outcome influenced by parental education and income

December 21 2020

---



Credit: CC0 Public Domain

Children from educationally deprived, low-income families are much more likely to face long-term cognitive impairment following a stroke. Such is the main finding of a recent study conducted at MedUni Vienna's Department of Pediatrics and Adolescent Medicine under the supervision of Lisa Bartha-Doering and Rainer Seidl.

With an incidence of between two and five cases per 100,000 [children](#), stroke in childhood is a rare occurrence. Nonetheless, this still means that up to 75 children in Austria suffer a stroke every year. Nearly half of these are left with long-term neurological impairment following a stroke. Apart from physical and motor deficits, they are often left with impaired speech, attention span, working speed, perception and memory. The working group led by Lisa Bartha-Doering and Rainer Seidl has now shown that children from educationally deprived and [low-income families](#) are more likely to suffer [long-term cognitive impairment](#) following a stroke.

Says Lisa Bartha-Doering: "Not all children have an equal chance of growing up healthy. The correlation between socioeconomic background, which encompasses the social and economic circumstances of the children, and their health has been documented for many years now. The recognition that socioeconomic status affects the outcome following a neurological disease is therefore not new but it is important to keep reminding ourselves of it. However, what did surprise us was the extent to which the variation in the cognitive outcome for children following a stroke can be explained by their [socioeconomic background](#)."

In fact, the study showed that nearly half of the differences in cognitive abilities were attributable to the children's socioeconomic status. Rainer Seidl, Head of the neuropediatric out-patient clinic in Vienna General Hospital/MedUni Vienna explains: "The opportunity for early rehabilitation after the acute phase has improved over the last few years,

due to the creation of children's rehabilitation centers and children's access to rehabilitation. However, the financial status of their families continues to influence access to important ongoing outpatient therapies, as well as their frequency and duration. Since therapists with statutory health insurance contracts often have long waiting lists, access to free physiotherapy, speech [therapy](#) and occupational therapy is limited in Austria. In rural regions, there is even less availability of free therapy places. Neurocognitive training is not funded at all by the health insurance fund in Austria. Higher-income parents therefore often finance the necessary treatment privately or pay for additional treatment sessions. Therefore, children from low-income families often cannot have the same number of treatment sessions as children of higher socioeconomic status."

The researchers also point to the cognitive reserve of the child (i.e. their individual flexibility for reorganizing cognitive processes in the brain) as another important factor. When researching the cognitive reserve, it is assumed that this is determined by the interplay of genetic and environmental factors: In healthy children, this reserve is closely linked to their intellectual achievements. The capacity for neuronal reorganization and its influence upon specific cognitive abilities play a major role, particularly following a stroke in childhood. The authors suppose a third important reason for the outcome of the study, which is the higher incidence of childhood strokes among educationally deprived and low-income classes. Although stroke in childhood and adolescence is less influenced by the typical adult risk factors such as diabetes, atherosclerosis, hypertension or smoking, low socioeconomic status is a risk factor for the incidence of stroke in childhood and its course, due to associated greater susceptibility to infection, poor diet and inadequately treated metabolic diseases.

"Our study shows that more attention needs to be given to the treatment of children from educationally deprived, low-income families. Free

access to treatment, including physiotherapy, speech therapy, occupational therapy and neurocognitive therapy should be available to all children who have suffered a [stroke](#). Teachers and caregivers should also be made aware of this particularly vulnerable patient group of children of lower [socioeconomic status](#), so that they can create a learning-friendly, supportive environment for them," conclude the authors in summary.

**More information:** Lisa Bartha-Doering et al. Influence of socioeconomic status on cognitive outcome after childhood arterial ischemic stroke, *Developmental Medicine & Child Neurology* (2020). [DOI: 10.1111/dmcn.14779](https://doi.org/10.1111/dmcn.14779)

Provided by Medical University of Vienna

Citation: Stroke in childhood and adolescence: Outcome influenced by parental education and income (2020, December 21) retrieved 5 May 2024 from <https://medicalxpress.com/news/2020-12-childhood-adolescence-outcome-parental-income.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--