


Participation in competitive sport in adolescence brings midlife health benefits to women

December 7 2020



Credit: CC0 Public Domain

Females who participate in competitive sport during adolescence have better fitness at midlife than do females with no competitive sport

background in adolescence, reveals a study conducted at the University of Jyväskylä. Higher lean mass and bone density and better physical performance at midlife were associated with competitive sport participation at the age of 13 to 16 years. The study also found that bone density was lower if the woman has had her first period at age 14 years or older.

The findings emphasize the link between adolescence [competitive sport](#) participation, and body composition, bone health and [physical performance](#) later in life.

"The relationship between adolescent participation in competitive sport and midlife health benefits were seen even if we took midlife physical activity into account," says Suvi Ravi, a Ph.D. student and the corresponding author at the Faculty of Sport and Health Sciences.

"We also investigated if individuals engaged in competitive sport in adolescence had more musculoskeletal problems than those with no competitive sport background in adolescence, but did not find any association between those factors," Ravi explains.

Another main finding of the study was that women whose period had occurred at age 14 years or older had [lower bone density](#) than women whose period had started at age 12 or younger. In women under menopausal age, this association was found even if physical activity in adolescence and at midlife was taken into account.

"It seems that a later age for a female's [first period](#) is associated with lower [bone density](#) regardless of competitive sport participation in adolescence," Ravi says. "It is known that some [female athletes](#) have their first period well above the average menarcheal age, so it is important to pay attention to girls whose menarche is delayed. Nowadays it is thought that menarche is belated if periods have not occurred by the

age of 15 years."

This study was part of the Estrogenic Regulation of Muscle Apoptosis (ERMA) study conducted at the University of Jyväskylä and Gerontology Research Center and led by Academy Research Fellow Eija Laakkonen. Nearly 1,100 (n = 1098) women between the ages of 47 and 55 participated in this part of the ERMA project where the associations between adolescence physical activity, age at menarche and midlife characteristics were investigated.

More information: Suvi Ravi et al, Adolescent Sport Participation and Age at Menarche in Relation to Midlife Body Composition, Bone Mineral Density, Fitness, and Physical Activity, *Journal of Clinical Medicine* (2020). [DOI: 10.3390/jcm9123797](https://doi.org/10.3390/jcm9123797)

Provided by University of Jyväskylä

Citation: Participation in competitive sport in adolescence brings midlife health benefits to women (2020, December 7) retrieved 23 April 2024 from <https://medicalxpress.com/news/2020-12-competitive-sport-adolescence-midlife-health.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.